




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> :  <b>C07D 487/04, 471/04, A61K 31/505, 31/435 // (C07D 487/04, 239:00, 231:00) (C07D 471/04, 249:00, 221:00) (C07D 487/04, 239:00, 209:00)</b></p>	<p><b>A1</b></p>	<p>(11) International Publication Number: <b>WO 98/08847</b></p> <p>(43) International Publication Date: <b>5 March 1998 (05.03.98)</b></p>
<p>(21) International Application Number: <b>PCT/IB97/00922</b></p> <p>(22) International Filing Date: <b>25 July 1997 (25.07.97)</b></p> <p>(30) Priority Data:  <b>60/025,039                      28 August 1996 (28.08.96)                      US</b></p> <p>(71) Applicant (for all designated States except US): <b>PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US).</b></p> <p>(72) Inventor; and  (75) Inventor/Applicant (for US only): <b>CHEN, Yuhpyng, Liang [US/US]; 8 Waterview Drive, Waterford, CT 06385 (US).</b></p> <p>(74) Agents: <b>SPIEGEL, Allen, J. et al.; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US).</b></p>		<p>(81) Designated States: <b>AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</b></p> <p><b>Published</b>  <i>With international search report.</i></p>

(54) Title: SUBSTITUTED 6,5-HETERO-BICYCLIC DERIVATIVES

## (57) Abstract

This invention relates to compounds of formula (I), wherein the dashed lines represent optional double bonds; A is nitrogen or CR<sup>7</sup>; B is -NR<sup>1</sup>R<sup>2</sup>, -CR<sup>1</sup>R<sup>2</sup>R<sup>10</sup>, C(-CR<sup>2</sup>R<sup>11</sup>)R<sup>1</sup>, -NHCR<sup>1</sup>R<sup>2</sup>R<sup>10</sup>, -OCR<sup>1</sup>R<sup>2</sup>R<sup>10</sup>, -SCR<sup>1</sup>R<sup>2</sup>R<sup>10</sup>, -CR<sup>2</sup>R<sup>10</sup>NHR<sup>1</sup>, -CR<sup>2</sup>R<sup>10</sup>OR<sup>1</sup>, -CR<sup>2</sup>R<sup>10</sup>SR<sup>1</sup> or -COR<sup>2</sup>; J and K are each independently nitrogen or carbon and both J and K are not nitrogens; D and E are each selected, independently, from nitrogen, CR<sup>4</sup>, C=O, C=S, sulfur, oxygen, CR<sup>4</sup>R<sup>6</sup> and NR<sup>8</sup>; G is nitrogen or carbon; and their use for the prevention or inhibition of a disorder that can be treated by antagonizing CRF.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KR	Kenya	NI	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

5                    SUBSTITUTED 6,5- HETERO-BICYCLIC DERIVATIVES                    Background of the invention

                    This invention relates to certain pharmaceutically active substituted 6,5-hetero-bicyclic derivatives, pharmaceutical compositions containing them and methods of administering them to subjects in need of their corticotropin releasing factor antagonist activity.

                    The substituted heterocyclic derivatives claimed in this case exhibit activity as corticotropin releasing factor (hormone) CRF (CRH) antagonists.

                    CRF antagonists are mentioned in U.S. Patents 4,605,642 and 5,063,245 referring to peptides and pyrazolinones, respectively. They are also referred to in the following: PCT Patent Application PCT/IB95/00439, which designates the United States and was filed on June 6, 1995 and published on December 14, 1995; PCT Patent Application PCT/IB95/00373, which designates the United States and was filed on May 18, 1995 and published on December 21, 1995; U.S. Patent Application 08/448,539, which was filed in the PCT on Nov. 12, 1993 and entered the U.S. national phase on June 14, 1995; PCT Patent Application WO 95/10506, which was filed on October 12, 1993 and published on April 20, 1995, and U.S. Patent Application 08/481,413, which was filed in the PCT on November 26, 1993 and entered the U.S. national phase on July 24, 1995; U.S. Patent Application 08/254,820, which was filed on April 19, 1995; Provisional U.S. Patent Application 60/008,396, which was filed on December 8, 1995; and Provisional U.S. Patent Application 60/006,333, which was filed on November 8, 1995. All the foregoing patent applications are incorporated herein by reference in their entireties.

                    The importance of CRF antagonists is set out in the literature, e.g., P. Black, Scientific American SCIENCE & MEDICINE, 1995, p. 16-25; T. Lovenberg et al., Current Pharmaceutical Design, 1995, 1, 305-316; and United States Patent 5,063,245, which is referred to above. A recent outline of the different activities possessed by CRF antagonists is found in M. J. Owens et al., Pharm. Rev., Vol. 43, pages 425 to 473 (1991), also incorporated herein by reference. Based on the research described in these two and other references, CRF antagonists are effective in the treatment of a wide range of stress-related illnesses, mood disorders such as depression, major depressive disorder, single episode depression, recurrent depression, child abuse induced depression, postpartum depression, dysthymia, bipolar disorders and cyclothymia;


-2-

chronic fatigue syndrome; eating disorders such as anorexia and bulimia nervosa; generalized anxiety disorder; panic disorder; phobias; obsessive-compulsive disorder, post-traumatic stress disorder, pain perception such as fibromyalgia; headache; gastrointestinal diseases; hemorrhagic stress; ulcers; stress-induced psychotic  
 5 episodes; fever; diarrhea; post-operative ileus, colonic hypersensitivity; irritable bowel syndrome; Crohn's disease; spastic colon; inflammatory disorders such as rheumatoid arthritis and osteoarthritis; pain; asthma; psoriasis; allergies; osteoporosis; premature birth; hypertension, congestive heart failure; sleep disorders; neurodegenerative diseases such as Alzheimer's disease, senile dementia of the Alzheimer's type,  
 10 multiinfarct dementia, Parkinson's disease, and Huntington's disease; head trauma; ischemic neuronal damage; excitotoxic neuronal damage; epilepsy; stroke; spinal cord trauma; psychosocial dwarfism; euthyroid sick syndrome; syndrome of inappropriate antidiarrhetic hormone; obesity; chemical dependencies and addictions; drug and alcohol withdrawal symptoms; infertility, cancer; infertility; muscular spasms; urinary  
 15 incontinence; hypoglycemia and immune dysfunctions including stress induced immune dysfunctions, immune suppression and human immunodeficiency virus infections; and stress-induced infections in humans and animals.

The compounds of this invention are also believed to be inhibitors of CRH binding protein and therefore useful in the treatment of disorders the treatment of which  
 20 can be effected or facilitated by inhibiting such protein. Examples of such disorders are Alzheimer's disease and obesity.

### Summary of the Invention

The present invention relates to compounds of the formula



or a pharmaceutically acceptable salt thereof, wherein

the dashed lines represent optional double bonds;

A is nitrogen or CR<sup>7</sup>;

-3-

B is  $-NR^1R^2$ ,  $-CR^1R^2R^{10}$ ,  $-C(=CR^2R^{11})R^1$ ,  $-NHCR^1R^2R^{10}$ ,  $-OCR^1R^2R^{10}$ ,  $-SCR^1R^2R^{10}$ ,  $-CR^2R^{10}NHR^1$ ,  $-CR^2R^{10}OR^1$ ,  $-CR^2R^{10}SR^1$  or  $-COR^2$ ;

J and K are each independently nitrogen or carbon and both J and K are not nitrogens ;

5 D and E are each selected, independently, from nitrogen,  $CR^4$ ,  $C=O$ ,  $C=S$ , sulfur, oxygen,  $CR^4R^6$  and  $NR^8$  ;

G is nitrogen or carbon;

the ring containing D, E, G, K, and J in formula I may be a saturated or unsaturated 5-membered ring and may optionally contain one or two double bonds and  
10 may optionally contain from one to three heteroatoms in the ring and may optionally have one or two  $C=O$  or  $C=S$  groups;

$R^1$  is  $C_1$ - $C_6$  alkyl optionally substituted with one or two substituents independently selected from hydroxy, fluoro, chloro, bromo, iodo,  $-O-(C_1-C_4 \text{ alkyl})$ ,  $CF_3$ ,  $-C(=O)O-(C_1-C_4 \text{ alkyl})$ ,  $-OC(=O)(C_1-C_4 \text{ alkyl})$ ,  $-OC(=O)N(C_1-C_4 \text{ alkyl})(C_1-C_2 \text{ alkyl})$ ,  
15  $-NHCO(C_1-C_4 \text{ alkyl})$ ,  $-COOH$ ,  $-COO(C_1-C_4 \text{ alkyl})$ ,  $-CONH(C_1-C_4 \text{ alkyl})$ ,  $-CON(C_1-C_4 \text{ alkyl})(C_1-C_2 \text{ alkyl})$ ,  $-S(C_1-C_4 \text{ alkyl})$ ,  $-CN$ ,  $-NO_2$ ,  $-SO(C_1-C_4 \text{ alkyl})$ ,  $-SO_2(C_1-C_4 \text{ alkyl})$ ,  $-SO_2NH(C_1-C_4 \text{ alkyl})$  and  $-SO_2N(C_1-C_4 \text{ alkyl})(C_1-C_2 \text{ alkyl})$ , wherein each of the  $C_1-C_4$  alkyl groups in the foregoing  $R^1$  groups may optionally contain one or two double or triple bonds;

20  $R^2$  is  $C_1$ - $C_{12}$  alkyl which may optionally contain from one to three double or triple bonds, aryl or  $(C_1-C_4 \text{ alkylene})\text{aryl}$ , wherein said aryl and the aryl moiety of said  $(C_1-C_4 \text{ alkylene})\text{aryl}$  is selected from phenyl, naphthyl, thienyl, benzothienyl, pyridyl, quinolyl, pyrazinyl, pyrimidinyl, imidazolyl, furanyl, benzofuranyl, benzothiazolyl, isothiazolyl, pyrazolyl, pyrrolyl, indolyl, pyrrolopyridyl, oxazolyl and benzoxazolyl;  $C_3$ - $C_8$  cycloalkyl  
25 or  $(C_1-C_6 \text{ alkylene})(C_3-C_8 \text{ cycloalkyl})$ , wherein one or two of the carbon atoms of said cycloalkyl and the 5 to 8 membered cycloalkyl moieties of said  $(C_1-C_6 \text{ alkylene})(C_3-C_8 \text{ cycloalkyl})$  may optionally and independently be replaced by an oxygen or sulfur atom or by  $NZ^2$  wherein  $Z^2$  is selected from hydrogen,  $C_1$ - $C_4$  alkyl, benzyl and  $C_1$ - $C_4$  alkanoyl, and wherein each of the foregoing  $R^2$  groups may optionally be substituted with from  
30 one to three substituents independently selected from chloro, fluoro, hydroxy and  $C_1$ - $C_4$  alkyl, or with one substituent selected from bromo, iodo,  $C_1$ - $C_6$  alkoxy,  $-OC(=O)(C_1-C_6 \text{ alkyl})$ ,  $-OC(=O)N(C_1-C_4 \text{ alkyl})(C_1-C_2 \text{ alkyl})$ ,  $-S(C_1-C_6 \text{ alkyl})$ , amino,  $-NH(C_1-C_2 \text{ alkyl})$ ,  $-N(C_1-C_2 \text{ alkyl})(C_1-C_4 \text{ alkyl})$ ,  $-N(C_1-C_4 \text{ alkyl})CO-(C_1-C_4 \text{ alkyl})$ ,  $-NHCO(C_1-C_4 \text{ alkyl})$ ,

-4-

-COOH, -COO(C<sub>1</sub>-C<sub>4</sub> alkyl), -CONH(C<sub>1</sub>-C<sub>4</sub> alkyl), -CON(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -SH, -CN, -NO<sub>2</sub>, -SO(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>NH(C<sub>1</sub>-C<sub>4</sub> alkyl) and -SO<sub>2</sub>N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl);

5 -NR<sup>1</sup>R<sup>2</sup> or -CR<sup>1</sup>R<sup>2</sup>R<sup>10</sup> may form a saturated 3 to 8 membered carbocyclic ring which may optionally contain from one to three double bonds and wherein one or two of the ring carbon atoms of such 5 to 8 membered rings may optionally and independently be replaced by an oxygen or sulfur atom or by NZ<sup>3</sup> wherein Z<sup>3</sup> is hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, benzyl or C<sub>1</sub>-C<sub>4</sub> alkanoyl;

10 R<sup>3</sup> is hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, -O(C<sub>1</sub>-C<sub>4</sub> alkyl), chloro, fluoro, bromo, iodo, (C<sub>1</sub>-C<sub>2</sub> alkylene)-O-(C<sub>1</sub>-C<sub>2</sub> alkyl), (C<sub>1</sub>-C<sub>2</sub> alkylene)-OH, or -S(C<sub>1</sub>-C<sub>4</sub> alkyl);

each R<sup>4</sup> is, independently, hydrogen, (C<sub>1</sub>-C<sub>6</sub> alkyl), fluoro, chloro, bromo, iodo, hydroxy, cyano, amino, (C<sub>1</sub>-C<sub>2</sub> alkylene)-OH, CF<sub>3</sub>, CH<sub>2</sub>SCH<sub>3</sub>, nitro, -O(C<sub>1</sub>-C<sub>4</sub> alkyl), -N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -S(C<sub>1</sub>-C<sub>4</sub> alkyl), -CO(C<sub>1</sub>-C<sub>4</sub> alkyl), -C(=O)H or -C(=O)O(C<sub>1</sub>-C<sub>4</sub> alkyl);

15 R<sup>5</sup> is hydrogen, methyl or ethyl;

R<sup>6</sup> is hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl;

R<sup>5</sup> is phenyl, pyridyl, pyrazinyl, pyrimidyl, pyridazinyl and wherein each of the foregoing R<sup>5</sup> groups is substituted with from one to four substituents R<sup>13</sup> wherein one to three of said substituents may be selected, independently, from fluoro, chloro, C<sub>1</sub>-C<sub>6</sub> alkyl and -O(C<sub>1</sub>-C<sub>6</sub> alkyl) and one of said substituents may be selected from bromo, iodo, formyl, OH, (C<sub>1</sub>-C<sub>4</sub> alkylene)-OH, (C<sub>1</sub>-C<sub>4</sub> alkylene)-O-(C<sub>1</sub>-C<sub>2</sub> alkyl), -CN, -CF<sub>3</sub>, -NO<sub>2</sub>, -NH<sub>2</sub>, -NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -N(C<sub>1</sub>-C<sub>2</sub> alkyl)(C<sub>1</sub>-C<sub>6</sub> alkyl), -OCO(C<sub>1</sub>-C<sub>4</sub> alkyl), (C<sub>1</sub>-C<sub>4</sub> alkylene)-O-(C<sub>1</sub>-C<sub>4</sub> alkyl), -S(C<sub>1</sub>-C<sub>6</sub> alkyl), (C<sub>1</sub>-C<sub>4</sub> alkylene)-S-(C<sub>1</sub>-C<sub>4</sub> alkyl), -C(=O)O(C<sub>1</sub>-C<sub>4</sub> alkyl), -C(=O)(C<sub>1</sub>-C<sub>4</sub> alkyl), -COOH, -SO<sub>2</sub>NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>N(C<sub>1</sub>-C<sub>2</sub> alkyl)(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>NH<sub>2</sub>, -NHSO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl), -S(C<sub>1</sub>-C<sub>6</sub> alkyl) and -SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl), and wherein each of the C<sub>1</sub>-C<sub>4</sub> alkyl and C<sub>1</sub>-C<sub>6</sub> alkyl moieties in the foregoing R<sup>5</sup> groups may optionally have one or two double bonds;

20 R<sup>7</sup> is hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, halo (e.g., chloro, fluoro, iodo or bromo), hydroxy, -O(C<sub>1</sub>-C<sub>4</sub> alkyl), -C(=O)(C<sub>1</sub>-C<sub>4</sub> alkyl), -C(=O)O(C<sub>1</sub>-C<sub>4</sub> alkyl), -OCF<sub>3</sub>, -CF<sub>3</sub>, -CH<sub>2</sub>OH or 30 -CH<sub>2</sub>O(C<sub>1</sub>-C<sub>2</sub> alkyl);

R<sup>10</sup> is hydrogen, hydroxy, methoxy or fluoro;

R<sup>11</sup> is hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl; and

-5-

with the proviso that: a) when both J and K are carbon and D is CR<sup>4</sup> and E is nitrogen, then G can not be nitrogen; (b) when both J and K are carbons and D and G are nitrogens, then E can not be CR<sup>4</sup> or C=O or C=S; (c) when both J and K are carbons and D and E are carbons, then G can not be nitrogen; (d) when G is carbon, it must be double bonded to E; and (e) in the ring containing J, K, D, E and G, there can not be two double bonds adjacent to each other.

and the pharmaceutically acceptable salts of such compounds.

When the ring containing D, E, G, K and J is a 5-membered heteroaromatic ring, it may be, e.g., pyrazolo, imidazolo, thieno, furano, thiazolo, oxazolo, triazolo or thiadiazolo.


The term "alkyl", as used herein, unless otherwise indicated, includes saturated monovalent hydrocarbon radicals having straight branched or cyclic moieties or combinations thereof.

The term "alkoxy", as used herein, unless otherwise indicated, means -O-alkyl, where "alkyl" is defined as above.

Examples of more specific embodiments of formula I are the following, wherein (R)<sub>n</sub> represents from zero to two substituents, wherein such substituents are as defined above in the definition of formula I.


-6-

5




I - A

10




I - B

15




I - E

20



I - J

30




I - K




-7-

5




10

I - C




I - D

15




I - G




I - H

20



I - L

25




I - M

30


-8-

5




10

I - N




I - O

15




20

I - R




I - S

25



30


I - V



I - W


-9-

5




I - P

10




I - Q

15




I - T

20




I - U

25




I - X

30




I - B'


-10-



I-Y



I-Z



I-A'

25

More specific embodiments of this invention include compounds of the above formula I wherein B is  $-\text{CHR}^1\text{R}^2$ ,  $-\text{NR}^1\text{R}^2$ ,  $-\text{NHCHR}^1\text{R}^2$ ,  $-\text{OCHR}^1\text{R}^2$  or  $-\text{SCHR}^1\text{R}^2$ , and  $\text{R}^1$  is  $\text{C}_1$ - $\text{C}_6$  alkyl, which may optionally be substituted with one hydroxy, fluoro,  $\text{CF}_3$  or  $\text{C}_1$ - $\text{C}_4$  alkoxy group and which may optionally contain one double or triple bond; and  
 5  $\text{R}^2$  is benzyl or  $\text{C}_1$ - $\text{C}_6$  alkyl, which may optionally contain one double or triple bond, wherein said  $\text{C}_1$ - $\text{C}_6$  alkyl and the phenyl moiety of said benzyl may optionally be substituted with one fluoro,  $\text{CF}_3$ ,  $\text{C}_1$ - $\text{C}_2$  alkyl,  $\text{C}_1$ - $\text{C}_2$  alkoxy or chloro group.

Other more specific embodiments of this invention include compounds of formula I wherein  $\text{R}^3$  is methyl, ethyl, chloro or methoxy;  $\text{R}^4$  and  $\text{R}^6$  are selected from  
 10 hydrogen, methyl and ethyl;  $\text{R}^5$  is di- or tri-substituted phenyl, pyridyl, or pyrimidyl, in which the two or three substituents on said phenyl, pyridyl or pyrimidyl are selected, independently, from  $\text{C}_1$ - $\text{C}_4$  alkyl,  $-\text{O}(\text{C}_1\text{-}\text{C}_4\text{ alkyl})$ ,  $(\text{C}_1\text{-}\text{C}_2\text{ alkyl})\text{-O}(\text{C}_1\text{-}\text{C}_4\text{ alkyl})$ ,  $-\text{CF}_3$ ,  $-\text{OCF}_3$ ,  $-\text{CHO}$ ,  $-(\text{C}_1\text{-}\text{C}_4\text{ alkyl})\text{-OH}$ , cyano, chloro, fluoro, bromo and iodo, wherein each of the foregoing  $\text{C}_1$ - $\text{C}_4$  alkyl groups may optionally contain one double or triple bond;

15 Other more specific embodiments of this invention include compounds of the formula I wherein B is or contains an  $\text{NR}^1\text{R}^2$  or  $\text{CR}^1\text{R}^2\text{R}^{10}$  moiety which forms a saturated or unsaturated 5-membered carbocyclic ring wherein one of the ring carbons may optionally be replaced by a sulfur or oxygen atom.

Other more specific embodiments of this invention include compounds of the  
 20 formula I wherein A is nitrogen or  $\text{CR}^7$  and  $\text{R}^7$  is hydrogen or methyl.

Other more specific embodiments of this invention include compounds of the formula I wherein B is  $-\text{CR}^1\text{R}^2\text{R}^{10}$ ,  $-\text{C}(=\text{CR}^2\text{R}^{11})\text{R}^1$ ,  $-\text{OCR}^1\text{R}^2\text{R}^{10}$ ,  $-\text{SCR}^1\text{R}^2\text{R}^{10}$ ,  $-\text{CR}^2\text{R}^{10}\text{NHR}^1$ ,  $\text{CR}^2\text{R}^{10}\text{OR}^1$ ,  $-\text{CR}^2\text{R}^{10}\text{SR}^1$  or  $-\text{COR}^2$ .

Other more specific embodiments of this invention include compounds of the  
 25 formula I wherein A is CH, J and K are carbon and D, E, and G are nitrogen;

Other more specific embodiments of this invention include compounds of the formula I wherein J and D are nitrogen, K and G are carbon, and E is CH,  $\text{CCH}_3$ , or  $\text{CC}_2\text{H}_5$ .

Other more specific embodiments of this invention include compounds of the  
 30 formula I wherein B contains a  $-\text{NR}^1\text{R}^2$  or  $-\text{CR}^1\text{R}^2\text{R}^{10}$  moiety that is a saturated or unsaturated three to five membered ring.

Other more specific embodiments of this invention include compounds of the formula I wherein J and K are carbon, and  $\text{D-E-G}$  is  $\text{O-C}(\text{CH}_3)=\text{C}$ ,  $\text{O-CH}=\text{C}$ ,

-12-

S-C(CH<sub>3</sub>)=C, S-CH=C, N(CH<sub>3</sub>)-C(CH<sub>3</sub>)=C, N(CH<sub>3</sub>)-CH=C, O-N=C, S-N=C, N(CH<sub>3</sub>)-N=C, O-CH<sub>2</sub>N, S-CH<sub>2</sub>N, or NH-C(C<sub>1</sub>-C<sub>2</sub>alkyl)=C.

Examples of preferred compounds of the formula I:

- 7-(1-ethyl-propoxy)-5-methyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidine;  
 5 [2,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-7-yl]-(1-ethyl-propyl)-amine;  
 (1-ethyl-propyl)-[5-methyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-7-yl]-amine;  
 7-(1-ethyl-propoxy)-2,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidine;  
 10 dine;  
 [2,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-7-yl]-ethyl-propyl-amine;  
 [6-bromo-5-bromomethyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-b]-pyridin-7-yl]-(1-ethyl-propyl)-amine;  
 15 (1-ethyl-propyl)-[5-methyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-b]-pyridin-7-yl]-amine;  
 [6-bromo-5-methyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-7-yl]-(1-ethyl-propyl)-methyl-amine; and  
 7-(1-ethyl-propoxy)-5-methyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-b]-pyridine.  
 20

Other compounds of formula I include the following:

- 4-(1-ethyl-propoxy)-2,7-dimethyl-8-(2,4,6-trimethyl-phenyl)-pyrrolo[1,2-a]-pyrimidine;  
 (1-ethyl-propyl)-[5-methyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-7-yl]-amine;  
 25 [3-(4-bromo-2,6-dimethyl-phenyl)-5-methyl-pyrazolo[1,5-a]pyrimidin-7-yl]-(1-ethyl-propyl)-amine;  
 butyl-ethyl-[5-methyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-7-yl]-amine;  
 30 3-(4-bromo-2,6-dimethyl-phenyl)-7-(1-ethyl-propoxy)-5-methyl-pyrazolo[1,5-a]-pyrimidine;  
 [2,7-dimethyl-8-(2,4,6-trimethyl-phenyl)-pyrrolo[1,2-a]pyrimidin-4-yl]-diethyl-amine;  
 [2,7-dimethyl-8-(2,4,6-trimethyl-phenyl)-pyrrolo[1,2-a]pyrimidin-4-yl]-ethyl-propyl-

-13-

amine;

[8-(4-chloro-2,6-dimethyl-phenyl)-2,7-dimethyl-pyrrolo[1,2-a]pyrimidin-4-yl]-ethyl-propyl-amine;

[8-(4-chloro-2,6-dimethyl-phenyl)-2,7-dimethyl-pyrrolo[1,2-a]pyrimidin-4-yl]-

5 diethyl-amine;

[8-(4-chloro-2,6-dimethyl-phenyl)-2,6-dimethyl-pyrrolo[1,2-a]pyrimidin-4-yl]-

diethyl-amine;

[2,6-dimethyl-8-(2,4,6-trimethyl-phenyl)-pyrrolo[1,2-a]pyrimidin-4-yl]-diethyl-

amine;

10 [2,6-dimethyl-8-(2,4,6-trimethyl-phenyl)-pyrrolo[1,2-a]pyrimidin-4-yl]-ethyl-propyl-amine;

butyl-[2,6-dimethyl-8-(2,4,6-trimethyl-phenyl)-pyrrolo[1,2-a]pyrimidin-4-yl]-ethyl-amine;

[2,6-dimethyl-8-(2,4,6-trimethyl-phenyl)-pyrrolo[1,2-a]pyrimidin-4-yl]-(1-ethyl-

15 propyl)-amine;

[8-(4-chloro-2,6-dimethyl-phenyl)-2,6-dimethyl-pyrrolo[1,2-a]pyrimidin-4-yl]-(1-ethyl-propyl)-amine;

8-(4-chloro-2,6-dimethyl-phenyl)-4-(1-ethyl-propoxy)-2,6-dimethyl-pyrrolo[1,2-a]-pyrimidine;

20 4-(1-ethyl-propoxy)-2,6-dimethyl-8-(2,4,6-trimethyl-phenyl)-pyrrolo[1,2-a]-pyrimidine;

4-(1-ethyl-propoxy)-2-methyl-8-(2,4,6-trimethyl-phenyl)-pyrrolo[1,2-a]pyrimidine;

(1-ethyl-propyl)-[2-methyl-8-(2,4,6-trimethyl-phenyl)-pyrrolo[1,2-a]pyrimidin-4-yl]-amine;

25 butyl-ethyl-[2-methyl-8-(2,4,6-trimethyl-phenyl)-pyrrolo[1,2-a]pyrimidin-4-yl]-amine;

butyl-ethyl-[5-methyl-3-(2,4,6-trimethyl-phenyl)-2H-isothiazolo[2,3-a]-pyrimidin-7-yl]-amine;

(1-ethyl-propyl)-[5-methyl-3-(2,4,6-trimethyl-phenyl)-2H-isothiazolo[2,3-a]-

30 pyrimidin-7-yl]-amine;

7-(1-ethyl-propoxy)-5-methyl-3-(2,4,6-trimethyl-phenyl)-2H-isothiazolo[2,3-a]-pyrimidine;

4-(1-ethyl-propoxy)-2-methyl-8-(2,4,6-trimethyl-phenyl)-imidazo[1,5-a]-

pyrimidine;

(1-ethyl-propyl)-[2-methyl-8-(2,4,6-trimethyl-phenyl)-imidazo[1,5-a]pyrimidin-4-yl]-  
amine;

5 [2,6-dimethyl-8-(2,4,6-trimethyl-phenyl)-imidazo[1,5-a]pyrimidin-4-yl]-(1-ethyl-  
propyl)-amine;

4-(1-ethyl-propoxy)-2,6-dimethyl-8-(2,4,6-trimethyl-phenyl)-imidazo[1,5-a]-  
pyrimidine;

4-(1-ethyl-propoxy)-2,6-dimethyl-8-(2,4,6-trimethyl-phenyl)-6,7-dihydro-imidazo  
[1,5-a]pyrimidine;

10 4-(1-ethyl-propoxy)-2,6-dimethyl-8-(2,4,6-trimethyl-phenyl)-thiazolo[3,4-a]-  
pyrimidine;

4-(1-ethyl-propoxy)-2-methyl-7-(2,4,6-trimethyl-phenyl)-pyrrolo[1,2-b]pyridazine;

4-(1-ethyl-propoxy)-2,5-dimethyl-7-(2,4,6-trimethyl-phenyl)-pyrrolo[1,2-b]-  
pyridazine;

15 4-(1-ethyl-propoxy)-2,6-dimethyl-7-(2,4,6-trimethyl-phenyl)-pyrrolo[1,2-b];  
pyridazine;

4-(1-ethyl-propoxy)-2,5,6-trimethyl-7-(2,4,6-trimethyl-phenyl)-pyrrolo[1,2-b]-  
pyridazine;

20 [2,5-dimethyl-7-(2,4,6-trimethyl-phenyl)-pyrrolo[1,2-b]pyridazin-4-yl]-(1-ethyl-  
propyl)-amine;

(1-ethyl-propyl)-[2-methyl-7-(2,4,6-trimethyl-phenyl)-pyrrolo[1,2-b]pyridazin-4-yl]-  
amine;

[2,6-dimethyl-3-(2,4,6-trimethyl-phenyl)-imidazo[1,2-b]pyridazin-8-yl]-(1-ethyl-  
propyl)-amine;

25 8-(1-ethyl-propoxy)-2,6-dimethyl-3-(2,4,6-trimethyl-phenyl)-imidazo[1,2-b]-  
pyridazine;

8-(1-ethyl-propoxy)-6-methyl-3-(2,4,6-trimethyl-phenyl)-imidazo[1,2-b]pyridazine;

(1-ethyl-propyl)-[6-methyl-3-(2,4,6-trimethyl-phenyl)-imidazo[1,2-b]pyridazin-8-yl]-  
amine;

30 (1-ethyl-propyl)-[2-methyl-7-(2,4,6-trimethyl-phenyl)-imidazo[1,5-b]pyridazin-4-yl]-  
amine;

[2,5-dimethyl-7-(2,4,6-trimethyl-phenyl)-imidazo[1,5-b]pyridazin-4-yl]-(1-ethyl-  
propyl)-amine;



-15-

4-(1-ethyl-propoxy)-2,5-dimethyl-7-(2,4,6-trimethyl-phenyl)-imidazo[1,5-b]-pyridazine-b]pyridazine;

butyl-[2,5-dimethyl-7-(2,4,6-trimethyl-phenyl)-imidazo[1,5-b]pyridazin-4-yl]-ethyl-amine;

5 [2,5-dimethyl-7-(2,4,6-trimethyl-phenyl)-imidazo[1,5-b]pyridazin-4-yl]-ethyl-propyl-amine;

[2,5-dimethyl-7-(2,4,6-trimethyl-phenyl)-imidazo[1,5-b]pyridazin-4-yl]-diethyl-amine;

10 diethyl-[6-methyl-3-(2,4,6-trimethyl-phenyl)-[1,2,4]triazolo[4,3-b]pyridazin-8-yl]-amine;

ethyl-[6-methyl-3-(2,4,6-trimethyl-phenyl)-[1,2,4]triazolo[4,3-b]pyridazin-8-yl]-propyl-amine;

butyl-ethyl-[6-methyl-3-(2,4,6-trimethyl-phenyl)-[1,2,4]triazolo[4,3-b]pyridazin-8-yl]-amine;

15 (1-ethyl-propyl)-[6-methyl-3-(2,4,6-trimethyl-phenyl)-[1,2,4]triazolo[4,3-b]-pyridazin-8-yl]-amine;

8-(1-ethyl-propoxy)-6-methyl-3-(2,4,6-trimethyl-phenyl)-[1,2,4]triazolo[4,3-b]-pyridazine;

20 7-(1-ethyl-propoxy)-1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]-pyridine;

7-(1-ethyl-propoxy)-1,2,5-trimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]-pyridine;

(1-ethyl-propyl)-[1,2,5-trimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]-pyridin-7-yl]-amine;

25 [1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]pyridin-7-yl]-(1-ethyl-propyl)-amine;

butyl-[1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]pyridin-7-yl]-ethyl-amine;

30 [1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]pyridin-7-yl]-ethyl-propyl-amine;

[1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]pyridin-7-yl]-diethyl-amine;

ethyl-[5-methyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-7-yl]-

- propyl-amine;
- butyl-ethyl-[5-methyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-7-yl]-amine;
- diethyl-[5-methyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-7-yl]-
- 5 amine;
- [3-(4-bromo-2,6-dimethyl-phenyl)-5-methyl-3H-[1,2,3]triazolo[4,5-b]pyridin-7-yl]-
- (1-ethyl-propyl)-amine;
- [3-(4-chloro-2,6-dimethyl-phenyl)-5-methyl-3H-[1,2,3]triazolo[4,5-b]pyridin-7-yl]-
- (1-ethyl-propyl)-amine;
- 10 3-(4-chloro-2,6-dimethyl-phenyl)-7-(1-ethyl-propoxy)-5-methyl-3H-[1,2,3]triazolo[4,5-b]pyridine;
- 3-(4-bromo-2,6-dimethyl-phenyl)-7-(1-ethyl-propoxy)-5-methyl-3H-[1,2,3]triazolo[4,5-b]pyridine;
- 7-(1-ethyl-propoxy)-5-methyl-3-(2,4,6-trimethyl-phenyl)-isothiazolo[4,5-b]pyridine;
- 15 (1-ethyl-propyl)-[5-methyl-3-(2,4,6-trimethyl-phenyl)-isothiazolo[4,5-b]pyridin-7-yl]-amine;
- (1-ethyl-propyl)-[5-methyl-3-(2,4,6-trimethyl-phenyl)-isoxazolo[4,5-b]pyridin-7-yl]-amine;
- 7-(1-ethyl-propoxy)-5-methyl-3-(2,4,6-trimethyl-phenyl)-isoxazolo[4,5-b]pyridine;
- 20 7-(1-ethyl-propoxy)-5-methyl-3-(2,4,6-trimethyl-phenyl)-thieno[3,2-b]pyridine;
- 7-(1-ethyl-propoxy)-2,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-thieno[3,2-b]pyridine;
- [2,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-thieno[3,2-b]pyridin-7-yl]-(1-ethyl-propyl)-amine;
- (1-ethyl-propyl)-[5-methyl-3-(2,4,6-trimethyl-phenyl)-thieno[3,2-b]pyridin-7-yl]-
- 25 amine;
- (1-ethyl-propyl)-[5-methyl-3-(2,4,6-trimethyl-phenyl)-furo[3,2-b]pyridin-7-yl]-amine;
- [2,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-furo[3,2-b]pyridin-7-yl]-(1-ethyl-propyl)-amine;
- 7-(1-ethyl-propoxy)-2,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-furo[3,2-b]pyridine;
- 30 7-(1-ethyl-propoxy)-5-methyl-3-(2,4,6-trimethyl-phenyl)-furo[3,2-b]pyridine;
- 4-(1-ethyl-propoxy)-2,5-dimethyl-7-(2,4,6-trimethyl-phenyl)-6H-pyrrolo[3,4-b]-pyridine;
- 4-(1-ethyl-propoxy)-2,5,6-trimethyl-7-(2,4,6-trimethyl-phenyl)-6H-pyrrolo[3,4-b]-

pyridine;

(1-ethyl-propyl)-[2,5,6-trimethyl-7-(2,4,6-trimethyl-phenyl)-6H-pyrrolo[3,4-b]pyridin-4-yl]-amine;

5      [1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-2-thia-4-aza-inden-7-yl]-(1-ethyl-propyl)-amine;

7-(1-ethyl-propoxy)-1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-2-thia-4-aza-indene;

4-(1-ethyl-propoxy)-2,5-dimethyl-7-(2,4,6-trimethyl-phenyl)-furo[3,4-b]pyridine;

[2,5-dimethyl-7-(2,4,6-trimethyl-phenyl)-furo[3,4-b]pyridin-4-yl]-(1-ethyl-propyl)-amine;

10      [2,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-2H-pyrazolo[4,3-b]pyridin-7-yl]-(1-ethyl-propyl)-amine;

7-(1-ethyl-propoxy)-2,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-2H-pyrazolo[4,3-b]-pyridine;

7-(1-ethyl-propoxy)-5-methyl-3-(2,4,6-trimethyl-phenyl)-isothiazolo[4,3-b]pyridine;

15      (1-ethyl-propyl)-[5-methyl-3-(2,4,6-trimethyl-phenyl)-isothiazolo[4,3-b]pyridin-7-yl]-amine;

(1-ethyl-propyl)-[5-methyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-d]-pyrimidin-7-yl]-amine;

20      diethyl-[5-methyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-d]pyrimidin-7-yl]-amine;

ethyl-[5-methyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-d]pyrimidin-7-yl]-propyl-amine;

butyl-ethyl-[5-methyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-d]-pyrimidin-7-yl]-amine;

25      [2,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-7-yl]-diethyl-amine;

butyl-[2,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-7-yl]-ethyl-amine;

30      butyl-ethyl-[5-methyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-7-yl]-amine;

ethyl-[5-methyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-7-yl]-propyl-amine;

diethyl-[5-methyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-7-yl]-amine;

-18-

[3-(4-chloro-2,6-dimethyl-phenyl)-5-methyl-pyrazolo[1,5-a]pyrimidin-7-yl]-(1-ethyl-propyl)-amine;

[3-(4-chloro-2,6-dimethyl-phenyl)-5-methyl-pyrazolo[1,5-a]pyrimidin-7-yl]-(1-ethyl-propyl)-amine;

5        3-(4-chloro-2,6-dimethyl-phenyl)-7-(1-ethyl-propoxy)-5-methyl-pyrazolo[1,5-a]-pyrimidine;

8-(4-chloro-2,6-dimethyl-phenyl)-4-(1-ethyl-propoxy)-2-methyl-pyrazolo[1,5-a]-[1,3,5]triazine;

10        (1-ethyl-propyl)-[2-methyl-8-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a][1,3,5]triazin-4-yl]-amine;

4-(1-ethyl-propoxy)-2-methyl-8-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a][1,3,5]-triazine;

4-(1-ethyl-propoxy)-2,7-dimethyl-8-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a][1,3,5]-triazine;

15        [2,7-dimethyl-8-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a][1,3,5]triazin-4-yl]-(1-ethyl-propyl)-amine;

[1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrazolo[4,3-b]pyridin-7-yl]-(1-ethyl-propyl)-amine;

20        7-(1-ethyl-propoxy)-1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrazolo[4,3-b]-pyridine;

7-(1-ethyl-propoxy)-5-methyl-3-(2,4,6-trimethyl-phenyl)-isothiazolo[4,5-b]-pyridine;

(1-ethyl-propyl)-[5-methyl-3-(2,4,6-trimethyl-phenyl)-isothiazolo[4,5-b]pyridin-7-yl]-amine;

25        1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-7-(1-ethyl-propoxy)-1H-pyrrolo[3,2-b]pyridine;

1,5-dimethyl-3-(2,6-dimethyl-4-chloro-phenyl)-7-(1-ethyl-propoxy)-1H-pyrrolo[3,2-b]pyridine;

30        2,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-7-(1-ethyl-propoxy)-1H-pyrrolo[3,2-b]pyridine;

2,5-dimethyl-3-(2,6-dimethyl-4-chloro-phenyl)-7-(1-ethyl-propoxy)-1H-pyrrolo[3,2-b]pyridine;

7-(1-cyclopropylmethyl-propoxy)-1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]pyridine;

7-(1-cyclopropylmethyl-propoxy)-1,5-dimethyl-3-(2,6-dimethyl-4-chloro-phenyl)-1H-pyrrolo[3,2-b]pyridine;

5 1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-7-(tetrahydro-furan-3-yloxy)-1H-pyrrolo[3,2-b]pyridine;

1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-7-(S)-(tetrahydro-furan-3-yloxy)-1H-pyrrolo[3,2-b]pyridine;

10 1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-7-(1-propyl-butoxy)-1H-pyrrolo[3,2-b]pyridine;

7-sec-butylsulfanyl-1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]pyridine;

cyclopropylmethyl-[2,5-dimethyl-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo[3,2-b]pyrimidin-4-yl]-propyl-amine;

15 2-[1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]pyridin-7-ylamino]-butan-1-ol;

1,5-dimethyl-7-thiazolidin-3-yl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]pyridine;

[1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]pyridin-7-yl]-ethyl-(2,2,2-trifluoro-ethyl)-amine;

20 cyclopropylmethyl-[1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]pyridin-7-yl]-propyl-amine;

(1-ethyl-propyl)-[5-methyl-3-(2,4,6-trimethyl-phenyl)-isoxazolo[4,5-b]pyridin-7-yl]-amine; and

25 7-(1-ethyl-propoxy)-5-methyl-3-(2,4,6-trimethyl-phenyl)-isoxazolo[4,5-b]-pyridine.

This invention also relates to a pharmaceutical composition for the treatment, prevention or inhibition of (a) a disorder the treatment of which can be effected or facilitated by antagonizing CRF, including but not limited to disorders induced or facilitated by CRF, or (b) a disorder selected from inflammatory disorders such as  
30 rheumatoid arthritis and osteoarthritis, pain, asthma, psoriasis and allergies; generalized anxiety disorder; panic; phobias; obsessive-compulsive disorder; post-traumatic stress disorder; sleep disorders induced by stress; pain perception such as fibromyalgia; mood disorders such as depression, including major depression, single episode

-20-

depression, recurrent depression, child abuse induced depression, mood disorders associated with premenstrual syndrome, and postpartum depression; dysthemia; bipolar disorders; cyclothymia; chronic fatigue syndrome; stress-induced headache; cancer; irritable bowel syndrome, Crohn's disease; spastic colon; post operative ileus; ulcer; diarrhea; stress-induced fever; human immunodeficiency virus (HIV) infections; neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease and Huntington's disease; gastrointestinal diseases; eating disorders such as anorexia and bulimia nervosa; hemorrhagic stress; chemical dependencies and addictions (e.g., dependencies on alcohol, nicotine, cocaine, heroin, benzodiazepines, or other drugs); drug and alcohol withdrawal symptoms; stress-induced psychotic episodes; euthyroid sick syndrome; syndrome of inappropriate antidiarrhetic hormone (ADH); obesity; infertility; head traumas; spinal cord trauma; ischemic neuronal damage (e.g., cerebral ischemia such as cerebral hippocampal ischemia); excitotoxic neuronal damage; epilepsy; stroke; immune dysfunctions including stress induced immune dysfunctions (e.g., porcine stress syndrome, bovine shipping fever, equine paroxysmal fibrillation, and dysfunctions induced by confinement in chickens, sheering stress in sheep or human-animal interaction related stress in dogs); muscular spasms; urinary incontinence; senile dementia of the Alzheimer's type; multiinfarct dementia; amyotrophic lateral sclerosis; hypertension; tachycardia; congestive heart failure; osteoporosis; premature birth; and hypoglycemia in a mammal, including a human, comprising an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, that is effective in the treatment of such disorder, and a pharmaceutically acceptable carrier.

The invention also relates to a method for the treatment, prevention or inhibition of (a) a disorder the treatment of which can be effected or facilitated by antagonizing CRF, including but not limited to disorders induced or facilitator by CRF, or (b) a disorder selected from inflammatory disorders such as rheumatoid arthritis and osteoarthritis, pain, asthma, psoriasis and allergies; generalized anxiety disorder; panic; phobias; obsessive-compulsive disorder; post-traumatic stress disorder; sleep disorders induced by stress; pain perception such as fibromyalgia; mood disorders such as depression, including major depression, single episode depression, recurrent depression, child abuse induced depression, mood disorders associated with premenstrual syndrome, and postpartum depression; dysthemia; bipolar disorders;

-21-

cyclothymia; chronic fatigue syndrome; stress-induced headache; cancer; irritable bowel syndrome; Crohn's disease; spastic colon; post operative ileus; ulcer; diarrhea; stress-induced fever; human immunodeficiency virus (HIV) infections; neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease and Huntington's disease; 5 gastrointestinal diseases; eating disorders such as anorexia and bulimia nervosa; hemorrhagic stress; stress-induced psychotic episodes; euthyroid sick syndrome; syndrome of inappropriate antidiarrhetic hormone (ADH); obesity; infertility; head traumas; spinal cord trauma; ischemic neuronal damage (e.g., cerebral ischemia such as cerebral hippocampal ischemia); excitotoxic neuronal damage; epilepsy; stroke; 10 immune dysfunctions including stress induced immune dysfunctions (e.g., porcine stress syndrome, bovine shipping fever, equine paroxysmal fibrillation, and dysfunctions induced by confinement in chickens, sheering stress in sheep or human-animal interaction related stress in dogs); muscular spasms; urinary incontinence; senile dementia of the Alzheimer's type; multiinfarct dementia; amyotrophic lateral sclerosis; 15 chemical dependencies and addictions (e.g., dependencies on alcohol, nicotine, cocaine, heroin, benzodiazepines, or other drugs); drug and alcohol withdrawal symptoms; hypertension; tachycardia; congestive heart failure; osteoporosis; premature birth; and hypoglycemia in a mammal, including a human, comprising administering to a subject in need of said treatment an amount of a compound of the formula I, or a 20 pharmaceutically acceptable salt thereof, that is effective in treating such disorder.

This invention also relates to a method of treating or preventing a disorder or condition, the treatment or prevention of which can be effected or facilitated by inhibiting CRH binding protein, in a mammal, including a human, comprising administering to said mammal a CRH binding protein inhibiting amount of a compound 25 of the formula I, or a pharmaceutically acceptable salt thereof.


This invention also relates to a pharmaceutical composition for treating or preventing a disorder or condition, the treatment or prevention of which can be effected or facilitated by inhibiting CRH binding protein in a mammal, including a human, comprising a CRH binding protein inhibiting amount of a compound of the formula I, 30 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

This invention includes all optical isomers and other stereoisomers of compounds of the formula I. When such compounds contain one or more chiral


centers, it is understood that the invention includes the racemic mixtures as well as all individual enantiomers and diastereomers of such compounds, and mixtures thereof.

### Detailed Description of the Invention


5       The following compounds having the formulas II, III, IV and V are useful as intermediates in the synthesis of compounds of the formula I.




II



III



IV



V

wherein T is chloro, bromo, iodo or -OSO<sub>2</sub>CF<sub>3</sub>; W is cyano, formyl, or -COO(C<sub>0</sub>-C<sub>4</sub> alkyl)  
 30 and A, J, K, D, E, G, R<sup>3</sup>, and R<sup>5</sup> are as defined above with reference to formula I.


Compounds of the formula I may be prepared as described below. In the reaction schemes and discussion that follow, A, B, D, E, G, J, K, R<sup>3</sup>, R<sup>5</sup> and structural formulae I, II, III, IV, and V are defined as above.



-23-

SCHEME 1

5




10

VI-a


IV-a

-24-


## SCHEME 2





-25-

SCHEME 3


15

SCHEME 4


-26-

SCHEME 5SCHEME 6

-27-


SCHEME 7

-28-

SCHEME 8

-29-

## SCHEME 9



-30-

Compounds of the formula I wherein B is  $-NR^1R^2$  or  $-NHCR^1R^2R^{11}$  may be prepared by reacting a compound of the formula II wherein T is chloro, bromo, or iodo with a compound of the formula BH, in the presence of a base, with or without an organometallic compound such as  $Cu(I)X$ , wherein X is chloro, bromo or iodo, or an acid (such as p-TsOH (Ts=Tosyl) or another sterically hindered phenol) or an equivalent agent known to those of skill in the art. Suitable solvents for this reaction include DMSO, NMP, dimethylacetamide and THF. An excess of BH may be used as both the reagent and the base. Other bases such as potassium or sodium carbonate, a trialkylamine, a potassium or sodium ( $C_1$ - $C_4$  alkoxide) and sodium hydride may also be used. When  $R^7$  is an electron withdrawing group such as  $-COO(C_1$ - $C_4$ alkyl) or CN, the reaction generally is carried out at a temperature between about room temperature and about  $100^\circ C$ . When  $R^7$  is a non-electron withdrawing group, the reaction temperature can generally range from about  $50^\circ C$  to about  $270^\circ C$  and the pressure can generally range from about 4 psi to about 300 psi. A pressure reactor may be used.

Alternatively, the compounds of formula I may be prepared by reacting a compound of the formula II wherein T is bromo or iodo with 1 equivalent or an excess of BH and a base such as sodium or potassium carbonate or a sodium or potassium ( $C_1$ - $C_4$  alkoxide), in the presence of a palladium (II) or a palladium (0) catalyst such as  $Pd(OAc)_2$  or  $Pd(PPh_3)_4$ , together with a racemic or chiral phosphino agent such as 2,2-bis(diphenylphosphino)-1,1-binaphthyl (BINAP). Alternatively, premade  $Pd(II)(BINAP)$  may be used in an appropriate inert (i.e., inert with respect to the reaction at hand) solvent such as toluene, xylene, dioxane or sulfolane, at a temperature from about room temperature to about  $180^\circ C$ , preferably at about reflux temperature.

Compounds of the formula I wherein B is  $-OCR^1R^2R^{11}$ ,  $-SCR^1R^2R^{11}$ , or  $-NHCR^1R^2R^{11}$  may be prepared by reacting compounds in the formula II wherein T is chloro, bromo or iodo with a compound of the formula BH in the presence of a base which is capable of deprotonation of BH (e.g., sodium or potassium hydride, or an organometallic base such as sodium diisopropylamide, sodium bis(trimethylsilyl)amide, lithium diisopropylamide, lithium bis(trimethylsilyl)amide, a sodium  $C_1$ - $C_4$  alkoxide or n-butyllithium), in an inert organic solvent such as tetrahydrofuran, acetonitrile, dimethylsulfoxide, acetone, a  $C_2$ - $C_5$  alcohol, chloroform, benzene, xylene, toluene, N,N-dimethylformamide (DMF), methylene chloride, 1-methyl-2-pyrrolidinone or a mixture of



two or more of the above solvents (e.g., DMSO and THF), at a temperature from about 0°C to about 180°C, preferably from about 50°C to about 180°C.

Compounds of the formula I wherein B is  $-\text{CR}^1\text{R}^2\text{R}^{11}$ ,  $-\text{C}(\text{C}=\text{CR}^2\text{R}^{12})\text{R}^1$ ,  $-\text{CR}^2\text{R}^{11}\text{NHR}^1$ ,  $-\text{CR}^2\text{R}^{11}\text{OR}^1$ ,  $-\text{CR}^2\text{R}^{11}\text{SR}^1$ , or  $-\text{C}(\text{O})\text{R}^2$  may be prepared from compounds  
5 of the formula III wherein W is cyano, formyl or carboxy, as described below.

Reacting compounds of formula III wherein W is cyano with a Grignard reagent containing the group  $\text{R}^2$  in an inert solvent such as THF, glyme, ether or dioxane, will yield corresponding compounds of formula I wherein B is  $-\text{COR}^2$ . Further reaction of the compounds of formula I wherein B is  $\text{COR}^2$  with a Grignard reagent containing  $\text{R}^1$ ,  
10 using a solvent such as those referred to above, will yield the corresponding the compounds of formula I wherein B is  $-\text{CR}^1\text{R}^2\text{OH}$ . Reacting compounds of formula III wherein W is formyl with a Grignard reagent containing the group  $\text{R}^2$ , in an ethereal solvent such as those referred to above, will yield the corresponding compounds of formula I wherein B is  $-\text{CHR}^2\text{OH}$ .

Compounds of formula I wherein B is  $-\text{CR}^1\text{R}^2\text{R}^{11}$  or  $-\text{C}(\text{C}=\text{CR}^2\text{R}^{11})\text{R}^1$  may be prepared by conventional methods. Thus, reaction of a compound of the formula I wherein B is  $-\text{CR}^1\text{R}^2\text{OH}$ , (wherein  $\text{R}^{1'}$  and  $\text{R}^{2'}$  are defined as  $\text{R}^1$  and  $\text{R}^2$ , respectively, except that  $\text{R}^{1'}$  may not be  $\text{R}^1$  and  $\text{R}^{2'}$  may not be  $\text{R}^2$ ), with an acid such as concentrated sulfuric acid in acetic acid, or a Burgess inner salt such as  
20 (carboxysulfamoyl)triethylammonium hydroxide methyl ester, will yield a compound of the formula I wherein B is  $-\text{C}(\text{C}=\text{CR}^2\text{R}^{11})\text{R}^1$ . Hydrogenation of a compound of formula I wherein B is  $-\text{C}(\text{C}=\text{CR}^2\text{R}^{11})\text{R}^1$  using palladium on carbon (Pd/C) or a platinum dioxide catalyst in a  $(\text{C}_1\text{-C}_4)$ alkanol solvent will yield a compound of the formula I wherein B is  $-\text{CHR}^1\text{R}^2$ . Reaction of a compound of the formula I wherein B is  $-\text{CR}^1\text{R}^2\text{OH}$  with  
25 diethylaminosulfur trifluoride or triphenylphosphine/carbon tetrachloride in an inert solvent such as carbon tetrachloride will afford a compound of the formula I wherein B is  $-\text{CR}^1\text{R}^2\text{F}$  or  $-\text{CR}^1\text{R}^2\text{Cl}$ , respectively.

Reduction of a compound of formula I wherein B is  $-\text{COR}^2$  with sodium borohydride, in an inert solvent such as a  $(\text{C}_1\text{-C}_4)$ alkanol, will yield a compound of the  
30 formula I wherein B is  $-\text{CHR}^2\text{OH}$ . Alkylation of a compound of the formula I wherein B is  $-\text{CHR}^2\text{OH}$  with an alkyl halide (such as alkyl iodide) in the presence of a base such as sodium hydride (NaH) at room temperature, in an inert solvent such as toluene, THF,

-32-

dioxane or ether, will yield the corresponding compound of the formula I wherein B is  $-\text{CHR}^2\text{OR}^1$ .

Compounds of the formula I wherein B is  $-\text{CR}^2\text{R}^{10}\text{NHR}^1$  may be prepared by conventional methods such as reductive amination of the corresponding compounds of the formula I wherein B is  $-\text{C}(\text{O})\text{R}^2$  with an appropriate amine and reducing agent (such as sodium cyanoborohydride, sodium triacetoxyborohydride, or lithium aluminum tetrahydride) in an appropriate inert solvent such as a  $\text{C}_1$ - $\text{C}_4$  alkanol or acetic acid.

Conversion of compounds in formula I wherein B is  $-\text{C}(\text{O})\text{R}^2$  into compounds in formula I wherein B is  $-\text{C}(\text{S})\text{R}^2$  can be accomplished using standard methods well known in the art (e.g., using Lawesson's Reagent or diphosphorus pentasulfide ( $\text{P}_2\text{S}_5$ )). Reduction of compounds of the formula I wherein B is  $-\text{C}(\text{S})\text{R}^2$  with a reducing agent such as sodium borohydride or lithium aluminum tetrahydride gives the corresponding compounds of the formula I wherein B is  $-\text{CHR}^2\text{SH}$ . Alkylation of compounds of the formula I wherein B is  $-\text{CHR}^2\text{SH}$  with alkyl halide (such as alkyl iodide) in the presence of a base such as sodium hydride in an inert solvent such as THF, DMF or toluene at about room temperature will afford the corresponding compounds of the formula I wherein B is  $-\text{CHR}^2\text{SR}^1$ .

Compounds in formula II may be prepared from compounds of the formula IV or V as described below.

Compounds of formula II wherein T is chloro, bromo or iodo can be prepared by reacting compounds of the formula IV with from one equivalent to an excess of  $\text{POT}_3$  (wherein T is chloro, bromo or iodo) in the presence or absence of a di( $\text{C}_1$ - $\text{C}_4$  alkyl)aniline, preferably diethylaniline, with or without a solvent (such as dichloroethane, DMF, dimethylsulfoxide (DMSO) or acetamide), at a temperature from about room temperature to about  $180^\circ\text{C}$ , preferably from about  $100^\circ\text{C}$  to about  $150^\circ\text{C}$ . Alternatively, compounds of formula II wherein T is chloro, bromo or iodo can be prepared by reacting the corresponding compounds of formula II wherein T is  $-\text{OSO}_2\text{CF}_3$  with a sodium or potassium halide, in an appropriate inert solvent such as THF, sulfolane, DMSO, DMF or acetonitrile, at a temperature from about  $60^\circ\text{C}$  to about  $180^\circ\text{C}$ . Compounds of formula II wherein T is  $-\text{OSO}_2\text{CF}_3$  can be prepared by reacting compounds of formula IV with  $\text{Tf}_2\text{O}$  in the presence of a base such as triethylamine or pyridine, in an appropriate inert solvent such as THF, methylene chloride, dioxane,

-33-

ether or toluene, at an temperature from about 0°C to about 50°C, preferably from about 0°C to about room temperature.

Alternatively, compounds of formula II wherein T is chloro, bromo or iodo may be prepared by reacting compounds of formula V with a (C<sub>1</sub>-C<sub>4</sub> alkyl)-nitrite and Cu(I)T<sub>2</sub> (wherein T is chloro, bromo or iodo) in an appropriate inert solvent such as acetonitrile, acetone, methylene chloride, THF, dioxane, benzene, toluene, dichloroethane, DMF, DMSO or N-methylpyrrolidinone (NMP) at a temperature from about room temperature to about 150°C, preferably from about 40°C to about 100°C.

Compounds of the formula IV may be prepared by reacting the appropriate compounds of formula V with sodium nitrite (NaNO<sub>2</sub>) in an aqueous acid such as sulfuric acid, acetic acid or phosphoric acid, with or without an organic solvent, preferably in acetonitrile (CH<sub>3</sub>CN) or acetone.

Compounds of formula III wherein W is cyano can be prepared by reacting the corresponding compounds of formula II wherein T is chloro, bromo or iodo with potassium cyanide, copper cyanide, sodium cyanide or a di(C<sub>1</sub>-C<sub>4</sub>alkyl)aluminum cyanide in an appropriate inert solvent such as dimethylsulfoxide, DMF, toluene or xylene, at temperature from about room temperature to about 180°C, preferably from about 60°C to about 150°C, with or without Pd(II)OAc or Pd(0)(PPh<sub>3</sub>)<sub>4</sub>.

Compounds of formula III wherein W is -CHO or -COOH may be prepared by reacting compounds in formula II wherein T is bromo or iodo with an organolithium reagent such as t-BuLi, s-BuLi, or n-BuLi in an appropriate inert solvent such as THF, dioxane, ether, benzene or methylene chloride, at temperature from about -120°C to about room temperature, preferably from about -110°C to about -60°C, followed by quenching with an appropriate electrophile such as DMF or CO<sub>2</sub> (gas or dry ice), to give compounds of formula III wherein W is -CHO and -COOH, respectively.

It is understood that the general organic chemistry knowledge can be applied to change the steps of the reaction sequences described herein depending on the feasibility of the reaction. For example, a protecting group may be used at any stage of the various syntheses described above at which it is workable, or an ester group may be reduced to the corresponding C<sub>1</sub>-C<sub>4</sub> alkyl group at any convenient stage. Compounds of formulas I - XVIII, wherein R<sup>3</sup> is chloro, bromo, -COO(C<sub>1</sub>-C<sub>4</sub> alkyl) or -COOH, can be converted to the corresponding compounds wherein R<sup>3</sup> is (C<sub>1</sub>-C<sub>4</sub> alkyl), -O(C<sub>1</sub>-C<sub>4</sub> alkyl), fluoro, -S(C<sub>1</sub>-C<sub>4</sub> alkyl) at any convenient stage, as appropriate, in the

-34-

syntheses referred to above using methods described in the literature. Compounds of formula I - XVIII, wherein  $R^3$  is  $-O(C_1-C_4 \text{ alkyl})$  or  $-S(C_1-C_4 \text{ alkyl})$  can be prepared by reacting the corresponding compounds wherein  $R^3$  is chloro, bromo or iodo, with a nucleophile such as a  $C_1-C_4$  alkanol or a  $C_1-C_4$  alkanethiol, in the presence of an organic or inorganic base. Suitable bases for such a reaction include sodium and sodium hydride. Compounds of formulas I - XVIII wherein  $R^3$  is fluoro can be prepared by reacting the corresponding compounds wherein  $R^3$  is chloro with tetrabutylammonium fluoride in a suitable inert solvent such as DMSO, methylene chloride, or tetrahydrofuran. Tetrahydrofuran is preferred.

- 10 Reduction of an ester or carboxylic acid using lithium aluminum tetrahydride/aluminum trichloride ( $LiAlH_4/AlCl_3$ ) in an appropriate inert solvent such as THF, ether, or dioxane, at temperature from about  $100^\circ C$  to about room temperature, will afford the corresponding compound wherein  $R^3$  is  $CH_3$ . Conversion of compounds wherein B is  $-COOH$  to the corresponding compounds wherein B is  $-CO(C_1-C_3 \text{ alkyl})$  may be accomplished using standard alkylation methods well known in art. Reduction of compounds wherein B is  $-CO(C_1-C_3 \text{ alkyl})$  using standard methods well known in the art will afford the corresponding compounds wherein  $R^3$  is one of various ( $C_1-C_4 \text{ alkyl})$  derivatives.

- 20 Compounds of the formula IV-a wherein A is  $CR^7$  and G, J, and K are carbon may be prepared by heating compounds of formula VI-a with an appropriate compound of the formula  $R^3C(O)CR^7COO(C_1-C_4 \text{ alkyl})$  under acid or Lewis acid conditions with or without a solvent, as shown in Scheme 1. Examples of such reaction conditions are: a) heating in polyphosphoric acid; b) heating in toluene, benzene or xylene in the presence of acid catalyst (such as p-TsOH, sulfuric acid or  $HCl(g)$ ) using a Dean-Stark trap; or c) heating in an appropriate solvent such as dichloroethane, diphenylether ( $Ph_2O$ ) or Dowtherm A in the presence of a Lewis acid such as  $SnCl_4$ ,  $ZnCl_2/HCl$  or  $AlCl_3$ .

- 30 Compounds of formula VI-a may be prepared using methods described in the scientific literature. (See Gazz. Chim. Ital.; 111, p167-172(1981); Chem. Pharm. Bull.; 24, 3001-3010 (1976); J. Org. Chem., 38, 1777-1780 (1973); Chem. Abstr. 68, 68982f (1968); Aust. J. Chem.; 22, 563-572(1969); J. Chem. Soc. Perkin Trans.2, p1954, p1957 (1972); J. Heterocycl. Chem.; FR, 19, 443-445(1982); J. Heterocycl. Chem., 22,

1496-1502 (1985); Tetrahedron, 47, 4639-4644 (1991); J. Heterocycl. Chem., 28, 2053-2055 (1991); J. Heterocycl. Chem., 29, 251-153 (1992).

Compounds of formula V-a and IV-a', wherein R is C<sub>1</sub>-C<sub>4</sub>alkyl can be prepared by heating compounds of formula VI wherein W is -CN and -COO(C<sub>1</sub>-C<sub>4</sub> alkyl), respectively, with an appropriate R<sup>3</sup>C(O)CH<sub>2</sub>COO(C<sub>1</sub>-C<sub>4</sub> alkyl) in the presence of a Lewis acid such as SnCl<sub>4</sub>, AlCl<sub>3</sub>, TiCl<sub>3</sub> or ZnCl<sub>2</sub>, in a inert solvent such as dichloroethane, at about the reflux temperature, as shown in Scheme 2. Base hydrolysis of a compound of the formula V-a or IV-a' with sodium hydroxide in H<sub>2</sub>O/(C<sub>1</sub>-C<sub>4</sub> alcohol) at reflux, or with lithium hydroxide in water/THF or water/dioxane at temperature from about room temperature to about the reflux temperature, followed by decarboxylation by heating in a oil bath at a temperature from about 140-180°C will give a compound of formula V-b or IV-b, respectively. Compounds of the formulas V-a and IV-a' may be converted into compounds of the formula II-a, and compounds of the formulas V-b and IV-b may be converted into compounds of the formula II-b by the procedures described above. The conversion of compounds of the formula II-a into those of the formula II-c can be accomplished using procedures analogous to those described above for converting compounds wherein R<sup>3</sup> is -COO(C<sub>1</sub>-C<sub>4</sub> alkyl) into those wherein R<sup>3</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl.

Compounds of formula VI may be prepared using methods described in the literature. (See Liebigs Ann Chem., 1534-1546, 1979; Gazz. Chim. Ital., 97, 25-33, 1967; Gazz. Chim. Ital., 120, 725-730, 1990; Eur. J. Med. Chem. Chim. Ther., 26, 143-158, 1991; J. Heterocycl. Chem., Fr. 19, 443-445, 1982; J. Heterocycl. Chem., 22, 1496-1502, 1985; J. Heterocycl. Chem., 31, 305-312, 1994; J. Heterocycl. Chem., 24, 243-245, 1987; J. Org. Chem. 57, 3713-3716, 1992; Liebigs Ann Chem., 1702-1710, 1984; Chem. Pharm. Bull., 34, 701-702, 1986; Pharmazie, 48, 849-853, 1993; Bull. Soc. Chim. Belg., 103, 181-184, 1994; and Indian. J. Chem. Sect. B, 33, 436-440, 1994).

Compounds of formula IV-c wherein A is N and G, J, and K are carbon may be prepared, as shown in Scheme 3, by reacting compounds of formula VI with (R<sup>3</sup>CO)<sub>2</sub>O, R<sup>3</sup>COOH or R<sup>3</sup>CO(OC<sub>1</sub>-C<sub>2</sub> alkyl)<sub>3</sub>, in acetic acid or an appropriate inert organic solvent such as toluene, dioxane, acetonitrile, methylene chloride or chloroform, at a temperature from about 25°C to about 150°C, preferably at about the reflux temperature, followed by heating in 85% phosphoric acid or an aqueous acid such as acetic acid, hydrochloric acid or sulfuric acid, preferably in 50-85% phosphoric acid. Compounds of formula V-b may be prepared, as shown in Scheme 3, by heating the

corresponding compounds of formula VI with excess of the appropriate compound of the formula  $R^3CONH_2$ .

Compounds of formula IV-d may be prepared, as shown in Scheme 4, by reacting compounds of formula IX with an appropriate reagent having the formula  $R^3C(O)CHR^7COO(C_1-C_4\text{alkyl})$  in an  $R^3COOH$  solvent at temperature from about 60°C to about 180°C, preferably at about the reflux temperature.

Compounds of formula I-N may be prepared, as shown in Scheme 5, by reacting the corresponding compounds of formula X with a  $(C_1-C_7\text{ alkyl})\text{nitrite}$ , with or without  $CuBr_2$ ,  $CuCl_2$ , or  $CuI_2$ , in an appropriate inert solvent such as acetonitrile, acetone, methylene chloride, chloroform, benzene or toluene, preferably acetonitrile at a temperature from about 25°C to about 150°C, preferably from about 60°C to 100°C.

Compounds of formula I-L and I-M may be prepared, as shown in Scheme 6, by reacting compounds of formula XI wherein X is S or O with a compound of the formula  $R^4CHO$  or  $R^4CH(OC_1-C_2\text{ alkyl})_2$  and an acid catalyst such as p-TsOH, HCl, HBr,  $H_2SO_4$ , or HCl, in toluene, xylene or benzene, preferably toluene, with from one to ten equivalents of water, at temperature from about 70°C to about 160°C, using a Dean-Stark trap or in the presence of anhydrous sodium sulfate.

Compounds of formula I-K and I-U may be prepared by reacting the corresponding compounds of formula XI with triphosgene and thiophosgene, respectively, and a base such as triethylamine or pyridine in an appropriate inert solvent such as methylene chloride, THF, dioxane, ether, benzene or chloroform, preferably methylene chloride or dry THF, at temperature from about 0°C to about 25°C.

Compounds of formula I-V, I-W, and I-X may be prepared, as illustrated in Scheme 7, starting from compounds of formula XII. Compounds of formula XIII may be prepared by reacting the corresponding compounds of formula XII with hydroxylamine in acidic (e.g., in trifluoroacetic acid) or basic (e.g., in NaOAc or NaOH and hydroxylamine hydrochloride in a  $C_1-C_4$  alcohol/water mixture) conditions at temperatures from about 25°C to about 150°C, preferably at about the reflux temperature. Compounds of formula XIV can be prepared by heating the corresponding compounds of formula XIII in acetic anhydride, trifluoroacetic anhydride or  $Tf_2O$ , with or without a solvent such as acetic acid or methylene chloride, in the presence of an appropriate amine base such as triethylamine or pyridine. Compounds of formula I-V, I-W and I-X may be prepared by heating the corresponding compounds

of formula XIV and pyridine in an inert solvent such as DMF, DMSO, NMP, sulfolane or acetamide at a temperature from about 80°C to about 180°C.

Compounds of formula I-G may be prepared, as illustrated in Scheme 8, by reducing the corresponding compounds of formula XVII. This reduction can be performed using standard methods known in literature for reduction of a nitro group to an amino group. Such methods include hydrogenation or reduction by iron in acetic acid. Cyclization may occur upon reduction or heating in an appropriate solvent such as a C<sub>1</sub>-C<sub>4</sub> alcohol, acetonitrile, toluene, THF, methylene chloride or acetic acid.

The conversion of compounds of formula XVI into those of formula XVII can be accomplished using methods analogous to those described above for the conversion of compounds of formula V wherein W is cyano into compounds of formula I wherein B is a group having a carbon atom directly attached to the bicyclic ring. The best method for conversion of a cyano group to a -COOH group is acid hydrolysis, for example, heating the cyano compound in 50-85% phosphoric acid or 50-90% acetic acid, preferably phosphoric acid. The best method for converting a cyano group into a -CO(C<sub>1</sub>-C<sub>4</sub> alkyl) is reacting the cyano compound with a Grignard reagent at a temperature from about 0°C to about 25°C in ether, THF or dioxane. The best method for converting a cyano group into a -CHO group is a diisobutylaluminum hydride reduction in THF, dioxane, or ether at a temperature from about -78°C to about 25°C, preferably from about -78°C to about -40°C.

Compounds of the formula XVI may be prepared by reacting compounds of formula XV, wherein Hal is chloro, bromo or iodo, with a sodium, potassium, or lithium salt of R<sup>5</sup>CH<sub>2</sub>CN in an appropriate inert solvent such as toluene, benzene, a C<sub>1</sub>-C<sub>5</sub> alcohol, THF, DMSO, dioxane, or pyridine, with or without a Pd(II) or Pd(0) catalyst, at a temperature from about -78°C to about 130°C.

Alternatively, compounds of formula XVII may be prepared by subjecting compounds of the formula XV to halogen-metal exchange (e.g., using an organo lithium agent such as tBuLi, s-BuLi or BuLi at -78°C in ether, THF, or dioxane), followed by quenching with an electrophile such as R<sup>5</sup>CHO, to give compounds of formula XVIII. Compounds of formula XIX-a may be prepared by reacting the corresponding compounds of formula XVIII with thionyl chloride, followed by reduction. Reacting compounds of formula XIX-a with a base (such as organolithium agent (e.g., lithium diisopropylamide or BuLi), followed by quenching with an electrophile (such as carbon

-38-

dioxide) will yield the corresponding compounds of formula XIX-b wherein  $R^{21}$  is -COOH. Compounds of formula XIX-c wherein  $R^{21}$  is -C(OH)R<sup>4</sup> may be prepared in a similar fashion by quenching the appropriate compound of formula XIX-a with an electrophile of the formula C<sub>1</sub>-C<sub>3</sub> alkyl-CHO. Compounds of formula XVII can be  
5 prepared by reacting compounds of formula XIX-c with PCC (pyridinium chlorochromate) using standard PCC oxidation methods that are well known in the art.

Compounds of formula II-d, II-e and II-f, wherein T is chloro, bromo or iodo and D, E, R<sup>3</sup>, R<sup>5</sup> and R<sup>7</sup> are as defined to those in formula I may be prepared, as shown in Scheme 9, by methods analogous to those described in Scheme 2. Alternatively,  
10 Compounds of formula II-d to II-f may be prepared using procedures analogous to those described in the literature (See: Pharm. Bull, 5, 229-231, 1957; Monatsh Chem, 100, 671-678, 1969; J. Heterocycl. Chem., 3, 218-220, 1966, Chem. Pharm. Bull., 23, 2891-2895, 1975; J. Heterocycl. Chem., 8, 1-6, 1971; Justus Liebigs Ann. Chem, 735, 35-44, 1970; Tetrahedron Lett, 1479, 1968; and Chem. Abstr., 1939, 4988.)

15 The acid addition salts of compounds of the formula can be prepared in a conventional manner by treating a solution or suspension of the corresponding free base with one chemical equivalent of a pharmaceutically acceptable acid. Conventional concentration or crystallization techniques can be employed to isolate the salts. Illustrative of suitable acids are acetic, lactic, succinic, maleic, tartaric, citric, gluconic,  
20 ascorbic, benzoic, cinnamic, fumaric, sulfuric, phosphoric, hydrochloric, hydrobromic, hydroiodic, sulfamic, sulfonic acids such as methanesulfonic, benzene sulfonic, p-toluenesulfonic, and related acids.

The compounds of formula I and their pharmaceutically acceptable salts (hereinafter referred to, collectively, as "the active compounds of this invention") may  
25 be administered alone or in combination with pharmaceutically acceptable carriers, in either single or multiple doses. Suitable pharmaceutical carriers include inert solid diluents or fillers, sterile aqueous solutions, oils (e.g., sesame oil, peanut oil) and various organic solvents. The pharmaceutical compositions formed by combining the novel compounds of formula I and pharmaceutically acceptable carriers can then be  
30 readily administered in a variety of dosage forms such as tablets, oil gel, powders, lozenges, syrups, injectable solutions and the like. These pharmaceutical compositions can, if desired, contain additional ingredients such as flavorings, binders, excipients and the like. Thus, for purposes of oral administration, tablets containing various excipients



such as sodium citrate, calcium carbonate and calcium phosphate may be employed along with various disintegrants such as starch, methylcellulose, alginic acid and certain complex silicates, together with binding agents such as polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in soft and hard filled gelatin capsules. Preferred materials for this include lactose or milk sugar and high molecular weight polyethylene glycols. When aqueous suspensions or elixirs are desired for oral administration, the essential active ingredient therein may be combined with various sweetening or flavoring agents, coloring matter or dyes and, if desired, emulsifying or suspending agents, together with diluents such as water, ethanol, propylene glycol, glycerin and combinations thereof.

For parenteral administration, solutions containing an active compound of this invention or a pharmaceutically acceptable salt thereof in sesame or peanut oil, aqueous propylene glycol, or in sterile aqueous solution may be employed. Such aqueous solutions should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. The sterile aqueous media employed are all readily available by standard techniques known to those skilled in the art.

The effective dosages for the active compounds of this invention will depend on the intended route of administration and factors such as the age and weight of the patient, as generally known to a physician. The dosages will also depend on the particular illness to be treated. For instance, the daily dosage for stress-induced illnesses, inflammatory disorders, Alzheimer's disease, gastro-intestinal diseases, anorexia nervosa, hemorrhagic stress and drug and alcohol withdrawal symptoms will generally range from about 0.1 to about 50 mg/kg body weight of the patient to be treated.

Methods that may be used to determine the CRF antagonist activity of the active compounds of this invention and their pharmaceutically acceptable salts are described in Endocrinology, 116, 1653-1659 (1985) and Peptides, 10, 179-188 (1985). The binding activities for compounds of the formula I, expressed as  $IC_{50}$  values, generally range from about 0.5 nanomolar to about 10 micromolar. Methods that can be used

-40-

to determine the CRF binding protein inhibiting activity of compounds of the formula I are described in Brain Research, (1997), 745 (1,2),248-255.

The present invention is illustrated by the following examples. It will be understood, however, that the invention is not limited to the specific details of these examples. Melting points are uncorrected. Proton nuclear magnetic resonance spectra ( $^1\text{H}$  NMR) and  $\text{C}^{13}$  nuclear magnetic resonance spectra ( $\text{C}^{13}$  NMR) were measured for solutions in deuteriochloroform ( $\text{CDCl}_3$ ) and peak positions are expressed in parts per million (ppm) downfield from tetramethylsilane (TMS). The peak shapes are denoted as follows: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; b, broad.

The following abbreviations are used in the Examples: Ph=phenyl; iPr=isopropyl.

#### EXAMPLE 1

##### 7-(1-Ethyl-propoxy)-5-methyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidine

A solution of 3-pentanol (140 mg, 1.5 mmol) in 1 ml of dry THF was treated with 60% sodium hydride in oil (28 mg, 0.7 mmol) and stirred at room temperature for 10 min. A solution of 7-chloro-5-methyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidine (75 mg, 0.262 mmol) in 1 ml of dry THF was added and the resulting mixture was heated at reflux for 5 hours. The mixture was quenched with water and extracted with ethyl acetate. The organic layer was washed with brine, dried and concentrated to give the crude material. The residue was purified through silica gel column chromatography using 3:7 chloroform:hexane as eluent to give 75mg (88%) of the title compound.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.97(s,1H), 6.97(s,2H), 6.03(s,1H), 4.56(m,1H), 2.53(s,3H), 2.32(s,3H), 2.13(s,6H), 2.10(m,4H), 1.09(t,6H) ppm.

#### EXAMPLE 2

##### [2,5-Dimethyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-7-yl]-(1-ethyl-propyl)-amine

A mixture of 7-Chloro-2,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidine (60 mg, 0.2 mmol) and 1-ethylpropylamine (4 ml) in 1ml of N-methylpyrrolidinone was heated at 125°C oil bath for 15 hrs. The mixture was quenched with water and extracted with ethyl acetate. The organic layer was dried and concentrated and purified through silica gel column chromatography using 20% ethyl acetate in hexane as eluent to give 35 mg of the title compound as a yellow solid.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$

-41-

6.96(s,2H), 6.00(d,1H), 5.77(s,1H), 3.47 (m,1H), 2.43(s,3H), 2.32(s,3H), 2.22(s,3H), 2.05(s,6H), 1.5-1.9(m,4H), 1.04(t,6H) ppm.

### EXAMPLE 3

#### (1-Ethyl-propyl)-[5-methyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-

#### 7-yl]-amine

A mixture of 7-chloro-5-methyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidine (190 mg, 0.63 mmol) and 1-ethylpropylamine (4 ml) in 1ml of N-methylpyrrolidinone was heated at 125°C oil bath for 15 hours. The mixture was quenched with water and extracted with ethyl acetate. The organic layer was dried and concentrated and purified through silica gel column chromatography using chloroform to 2% methanol in chloroform as eluent to give 195 mg (87%) of the title compound as a green solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.87(s,1H), 6.97(s,2H), 6.12(d,1H), 5.85(s,1H), 3.52(m,1H), 2.48(s,3H), 2.09(s,3H), 2.16(s,6H), 1.6-1.9(m,4H), 1.05(t,6H)ppm.

### EXAMPLE 4

#### 7-(1-Ethyl-propoxy)-2,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidine

To a suspension of 60% Sodium hydride in oil (160 mg) in 4 ml of DMSO was added 3-pentanol (853 mg), and then 7-chloro-2,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidine (580 mg) at room temperature. The mixture was heated at 88°C overnight. The mixture was quenched with water and extracted with ethyl acetate. The organic layer was dried and concentrated and purified through silica gel column chromatography using 10% hexane in chloroform as eluent to give the title compound as an orange oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 6.96(s,2H), 5.95(s,1H), 4.52(m,1H), 2.48(s,3H), 2.33(s,3H), 2.27(s,3H), 2.03(s,6H), 1.75-2.00(m,4H), 1.08(t,6H)ppm.

### EXAMPLE 5

#### [2,5-Dimethyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-7-yl]-ethyl-propyl-amine

A mixture of 7-chloro-5-methyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidine (200 mg, 0.66 mmol) and N-propylethylamine (2 ml) in 1ml of N-methylpyrrolidinone was heated at 135°C oil bath for 4 hours. The mixture was quenched with water and extracted with ethyl acetate. The organic layer was dried and concentrated and purified through silica gel column chromatography using hexane to 10% ethyl acetate in hexane as eluent to give 150 mg of the title compound as a clear

-42-

green oil.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  6.95(s,2H), 5.80(s,1H), 3.85(q,2H), 3.67(dd,2H), 2.41(s,3H), 2.32(s,3H), 2.21(s,3H), 2.03(s,6H), 1.76(m,2H), 1.29(t,3H), 0.98(t,3H) ppm.

#### EXAMPLE 6

5 [6-Bromo-5-bromomethyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-7-yl]-(1-ethyl-propyl)-amine

To a solution of butyl nitrite (119 mg, 1.15 mmol) and  $\text{CuBr}_2$  (205 mg, 0.919 mmol) in 16 ml of acetonitrile was added N4-(1-ethyl-propyl)-6-methyl-N2-(2,4,6-trimethyl-phenyl)-pyridine-2,3,4-triamine (250 mg, 0.766 mmol). The resulting mixture was heated at  $65^\circ\text{C}$  for 1.5 hrs. The mixture was cooled to room temperature and 2N HCl (16 ml) was added. The mixture was neutralized with 2N NaOH and extracted with ethyl acetate. The organic layer was washed with brine, dried and concentrated to give a brown oil. The oil residue was purified through silica gel column chromatography using 1:1 hexane :ethyl acetate as eluent to give 61mg of [6-bromo-5-bromomethyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-b] pyridin-7-yl]-(1-ethyl-propyl)-amine as white crystals. Mp  $123\text{--}125^\circ\text{C}$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.06(s,2H), 5.53(d,1H), 5.22(m,1H), 4.67(s,2H), 2.39(s,3H), 1.96(s,6H), 1.6-1.9(m,4H), 1.06(t,6H)ppm and 103 mg of [6-bromo-5-methyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-7-yl]-(1-ethyl-propyl)-amine as white solid. Mp  $115\text{--}117^\circ\text{C}$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.04(s,2H), 5.36(d,1H), 5.21(m,1H), 2.64(s,3H), 2.38(s,3H), 1.95(s,6H), 1.6-1.9(m,4H), 1.05(t,6H)ppm.

#### EXAMPLE 7

25 -(1-Ethyl-propyl)-[5-methyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-7-yl]-amine

A mixture of butyl nitrite (119 mg, 1.15 mmol) and N4-(1-ethyl-propyl)-6-methyl-N2-(2,4,6-trimethyl-phenyl)-pyridine-2,3,4-triamine (250 mg, 0.766 mmol) in anhydrous acetonitrile (16 ml) was heated at  $65^\circ\text{C}$  for 2 hours. The mixture was cooled to room temperature and 2N HCl (16 ml) was added. The mixture was neutralized with 2N NaOH and extracted with ethyl acetate. The organic layer was washed with brine, dried and concentrated to give 250 mg of the crude product as a brown oil. The oil residue was purified through silica gel column using chloroform as eluent to give 201 mg of the title compound as golden yellow solid. Mp  $131\text{--}133^\circ\text{C}$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.022(s,2H), 6.20(s,1H), 5.44(d,1H), 3.65(m,1H), 2.50(s,3H), 2.36(s,3H), 1.96(s,6H), 1.5-1.8(m,4H), 1.03(t,6H)ppm.

EXAMPLE 8

[6-Bromo-5-methyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-7-yl]-(1-ethyl-propyl)-methyl-amine

To a solution of [6-bromo-5-methyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo  
5 [4,5-b]pyridin-7-yl]-(1-ethyl-propyl)-amine (50 mg, 0.12 mmol) in dry THF (1.5 ml) was  
added 2.5 M n-BuLi in hexane (0.14 ml) at -78°C. After stirring at -78°C for 10 min, 0.5  
ml of methyl iodide was added at that temperature then allowed to warm to room  
temperature and stirred for 15 min for 2 hours. The mixture was quenched with water  
and extracted with ethyl acetate. The organic layer was washed with brine, dried and  
10 concentrated to give 46 mg of brown oil. The residue was purified through silica gel  
column chromatography using 1:1 chloroform:hexane as eluent to give the title  
compound as a golden oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.04(s,2H), 4.35(m,1H), 3.32(s,3H),  
2.70(s,3H), 2.38(s,3H), 1.94(s,6H), 1.7-2.0(m,4H), 1.01(t,6H)ppm.

EXAMPLE 9

15 7-(1-Ethyl-propoxy)-5-methyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]  
triazolo[4,5-b]pyridine

A mixture of 4-(1-Ethyl-propoxy)-6-methyl-N2-(2,4,6-trimethyl-phenyl)-pyridine-  
2,3-diamine (50 mg, 0.153 mmol) and butyl nitrite (24 mg, 0.229 mmol) in 4 ml  
acetonitrile was heated at 65°C for 2 hours. An additional 0.13 ml of butyl nitrite was  
20 added and the resulting mixture was heated at 65°C for 2 hours. The mixture was  
quenched with water and extracted with ethyl acetate. The organic layer was dried and  
concentrated to give 58 mg of brown oil. The oil was purified through silica gel column  
chromatography using 5% ethyl acetate in hexane as eluent to give 46 mg (88%) of the  
title compound as a light yellow oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.04(s,2H), 6.60(s,1H),  
25 5.26(m,1H), 2.57(s,3H), 2.38(s,3H), 1.94(s,6H), 1.8-2.0(m, 4H), 1.07(t,6H) ppm.

EXAMPLE 10

4-(1-Ethyl-propoxy)-2,5-dimethyl-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo[3,2-d]py-  
rimidine

To a solution of 3-pentanol (0.09 ml, 0.883 mmol) in dry THF was added 60%  
30 NaH in oil (20 mg, 0.500 mmol) and stirred for 5 min. A solution of 4-chloro-2,5-  
dimethyl-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo[3,2-d]pyrimidine (50 mg, 0.166 mmol) in  
dry THF was added to the reaction mixture and the resulting mixture was heated at  
reflux for 2 hr. The mixture was quenched with water, extracted with ethyl acetate. The

-44-

organic layer was washed with brine, separated, dried and concentrated to dryness to give the title compound as a white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 6.92(s,3H), 5.43(m,1H), 4.02(s,3H), 2.56(s,3H), 2.29(s,3H), 2.08(s,6H), 1.80(m,4H), 0.99(t,6H).

The title compounds of Examples 11-14 were prepared by the method analogous to that described in Example 10 starting from of 4-chloro-2-methyl-5-substituted-7-(substituted-phenyl)-5H-pyrrolo[3,2-d]pyrimidine or 7-chloro-5-methyl-1-substituted-3-(substituted-phenyl)-1H-pyrrolo[3,2-b]pyridine and an appropriate alcohol or thiol and a base.

#### EXAMPLE 11

10 (+)-2,5-Dimethyl-4-(tetrahydro-furan-3-yloxy)-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo[3,2-d]pyrimidine.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 6.95(s,1H), 6.92(s,2H), 5.88(m,1H), 3.9-4.08(m,4H), 4.01(s,3H), 2.56(s,3H), 2.29(s,3H), 2.2-2.4(m,2H), 2.07(s,6H) ppm.

#### EXAMPLE 12

15 2,5-Dimethyl-4-(S)-(tetrahydro-furan-3-yloxy)-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo[3,2-d]pyrimidine.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 6.95(s,1H), 6.92(s,2H), 5.88(m,1H), 3.9-4.08(m,4H), 4.01(s,3H), 2.56(s,3H), 2.29(s,3H), 2.2-2.4(m,2H), 2.07(s,6H) ppm.

#### EXAMPLE 13

20 2,5-Dimethyl-4-(1-propyl-butoxy)-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo[3,2-d]pyrimidine.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 6.93(s,2H), 6.92(s,1H), 5.58(m,1H), 4.02(s,3H), 2.56(s,3H), 2.29(s,3H), 2.09(s,6H), 1.6-1.8(m,4H), 1.4-1.6(m,4H), 0.96(t,6H)ppm.

#### EXAMPLE 14

25 4-sec-Butylsulfanyl-2,5-dimethyl-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo[3,2-d]pyrimidine.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 6.97(s,1H), 6.94(s,2H), 4.34(m,1H), 4.13(s,3H), 2.63(s,3H), 2.30(s,3H), 2.07(s,6H), 1.7-1.9(m,2H), 1.48(d,3H), 1.09(t,3H) ppm.

30 The title compounds of Examples 15-18 were prepared using the following procedure.

#### PROCEDURE FOR EXAMPLES. 15 - 18

A mixture of 4-chloro-2-methyl-5-substituted-7-(substituted-phenyl)-5H-pyrrolo[3,2-d]pyrimidine or 7-bromo-1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-

-45-

pyrrolo[3,2-b]pyridine-6-carboxylic acid methyl ester (1 mmol) and an appropriate amine in DMSO (2 ml) was heated in 130°C oil bath until all the starting material was consumed. The mixture was quenched with water and extracted with ethyl acetate. The organic layer was dried and concentrated to dryness to give the corresponding 4-alkylamino-2-methyl-5-substituted-7-(substituted-phenyl)-5H-pyrrolo[3,2-d]pyrimidin-  
 5 7-alkylamino-1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]pyridine-6-carboxylic acid methyl ester derivative. Silica gel column chromatography may be used for purification.

**EXAMPLE 15**

10 [2,5-Dimethyl-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo[3,2-d]pyrimidin-4-yl]-(1-ethyl-propyl)-amine

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 6.91(s,2H), 6.76(s,1H), 4.61(d,1H,NH), 4.33(m,1H), 4.04(s,3H), 2.49(s,3H), 2.28(s,3H), 2.09(s,6H), 1.72(m,2H), 1.60(m,2H), 0.98(t,6H)ppm.

**EXAMPLE 16**

15 Butyl-[2,5-dimethyl-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo[3,2-d]pyrimidin-4-yl]-ethyl-amine

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 6.96(s,1H), 6.92(s,2H), 3.93(s,3H), 3.44(q,2H), 3.40(m,2H), 2.57(s,3H), 2.29(s,3H), 2.09(s,6H), 1.57(m,2H), 1.30(m,2H), 1.14(t,3H), 0.88(t,3H)ppm.

**EXAMPLE 17**

20 2,5-Dimethyl-4-thiazolidin-3-yl-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo[3,2-d]pyrimidine

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.03(s,1H), 6.92(s,2H), 4.78(s,2H), 4.02(s,3H), 3.96(m,2H), 3.18(m,2H), 2.56(s,3H), 2.29(s,3H), 2.06(s,6H) ppm.

**EXAMPLE 18**

25 7-(1-Ethyl-propylamino)-1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]pyridine-6-carboxylic acid methyl ester

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 6.98(s,1H), 6.93(s,2H), 5.40(d,1H), 3.97(s,3H), 3.91(s,3H), 3.35(m,1H), 2.57(s,3H), 2.30(s,3H), 2.09(s,6H), 1.52(m,4H), 0.87(t,6H) ppm.

**EXAMPLE 19**

30 7-(1-Ethyl-propylamino)-1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]pyridine-6-carboxylic acid

A mixture of 7-(1-ethyl-propylamino)-1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]pyridine-6-carboxylic acid methyl ester and NaOH in a 1:1 mixture of

-46-

MeOH/water was heated at reflux over night. The resulting mixture was acidified with 2N HCl to a pH of 4-5, and extracted with chloroform. The organic layer was dried and concentrated to give the title compound.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.02(s,1H), 6.82(s,2H), 3.98(s,3H), 3.78(m,1H), 2.59(s,3H),  
5 2.07(s,3H), 2.00(s,6H), 1.64(m,4H), 0.90(t,6H) ppm.

#### EXAMPLE 20

##### 1,5-Dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]pyridin-7-yl)-(1-ethyl-propyl)-amine

A mixture of 7-(1-ethyl-propylamino)-1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-  
10 pyrrolo[3,2-b]pyridine-6-carboxylic acid was heated in 150-160°C oil bath until all the starting material was consumed. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 6.94(s,2H), 6.87(s,1H), 6.15(s,1H), 6.10(d,1H), 4.24(s,3H), 3.50(m,1H), 2.64(s,3H), 2.30(s,3H), 2.05(s,6H), 1.77(m,4H), 1.02(t,6H)ppm.

#### PREPARATION A

##### 15 4-(2,4,6-Trimethyl-phenyl)-2H-pyrazol-3-ylamine

A mixture of 3-oxo-2-(2,4,6-trimethyl-phenyl)-propionitrile (2.300 g, 12.3 mmol), hydrazine hydrate (0.93 g) and glacial acetic acid (1.55 ml) in 20 ml benzene was heated at reflux for 4.5 hours. Reaction mixture was cooled to room temperature and 50 ml of 18.5% HCl in water was added. The benzene layer was separated and  
20 reextracted with 18.5% HCl. The aqueous layer were combined and neutralized with ammonium hydroxide and stirred at rt. overnight. Precipitate formed and was filtered to yield the title compound (0.256 g) as a yellow solid. The benzene layer was concentrated and purified through silica gel column using 5% methanol in chloroform as eluent to give an additional 1.450 g of the title compound; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ  
25 7.24(s,1H), 6.95(s,2H), 4.75(brs,2H), 2.32(s,3H), 2.13(s,6H) ppm.

#### PREPARATION B

##### 5-Methyl-4-(2,4,6-trimethyl-phenyl)-2H-pyrazol-3-ylamine

The title compound was prepared as white solid by the method analogous to that described in preparation A starting from 3-oxo-2-(2,4,6-trimethyl-phenyl)-  
30 butyronitrile. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.7(brs,1H), 6.96(s,2H), 2.32(s,3H), 2.13(s,3H), 2.06(s,6H) ppm.



PREPARATION C2,5-Dimethyl-3-(2,4,6-trimethyl-phenyl)-4H-pyrazolo[1,5-a]pyrimidin-7-one

Methyl acetoacetate (0.38 ml) was added to a solution of 5-methyl-4-(2,4,6-trimethyl-phenyl)-2H-pyrazol-3-ylamine (641 mg, 2.98 mmol) in 4 ml of acetic acid. The reaction mixture was heated at reflux for overnight. The mixture was concentrated to dryness and the residue was purified through silica gel column chromatography using 5% methanol in chloroform as eluent to give 560 mg (65.5%) of the title compound as a white solid; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.7(s,1H), 6.98(s,2H), 5.52(s,1H), 2.29(s,3H), 2.20(s,3H), 1.97(s,3H), 1.95(s,6H) ppm.

PREPARATION D5-Methyl-3-(2,4,6-trimethyl-phenyl)-4H-pyrazolo[1,5-a]pyrimidin-7-one

Methyl acetoacetate (0.7 ml) was added to a solution of 4-(2,4,6-Trimethyl-phenyl)-2H-pyrazol-3-ylamine (1.120 g, 5.57 mmol) in 5 ml of acetic acid and the resulting mixture was heated at reflux for two days. Reaction mixture cooled and a white solid formed. Ethanol (6 ml) was added and stirred at room temperature overnight, filtered to give white solid which was recrystallized from ethanol to give 673 mg(45.2%) of the title compound as white crystals. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.9(s,1H), 7.7(s,1H), 6.95(s,2H), 5.55(s,1H), 2.25(s,3H), 2.20(s,3H), 2.0(s,6H)ppm.

PREPARATION E7-Chloro-5-methyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidine

A suspension of 5-methyl-3-(2,4,6-trimethyl-phenyl)-4H-pyrazolo[1,5-a]pyrimidin-7-one (590 mg, 2.2 mmol) in 9 ml of POCl<sub>3</sub> was treated with diethylaniline (0.7 ml) and the resulting mixture was stirred at reflux for 15 hours. The reaction mixture was concentrated to dryness. The residue was treated with ice water and stirred for 20 min, then extracted with chloroform. The organic layer was dried and concentrated to yield an orange oil which crystallized upon standing. The material was purified through silica gel column chromatography using chloroform as eluent to give 590 mg (94%) of the title compound as a yellow solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.08(s,1H), 6.98(s,2H), 6.86(s,1H), 2.56(s,3H), 2.33(s,3H), 2.09(s,6H) ppm.

### PREPARATION F

#### 7-Chloro-2,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidine

The title compound was prepared as an oil by the method analogous to that described in preparation E starting from 2,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-4H-pyrazolo[1,5-a]pyrimidin-7-one. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 6.98(s,2H), 6.77(s,1H), 2.51(s,3H), 2.33(s,3H), 2.31(s,3H), 1.99(s,6H) ppm.

### PREPARATION G

#### 2,5-Dimethyl-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo[3,2-d]pyrimidin-4-ol

A mixture of 3-amino-1-methyl-4-(2,4,6-trimethylphenyl)-1H-pyrrole-2-carbonitrile (0.4 mmol) and acetic anhydride (0.043 ml) in acetic acid (0.01 ml) was heated at reflux until all the starting material was consumed. The reaction mixture was concentrated to dryness. The residue was quenched with water and extracted with ethyl acetate. The organic extracts was washed with brine and concentrated to dryness. The residue was suspended in 0.5 ml of 85% phosphoric acid and heated at 130°C for 1 hour. The mixture was cooled and poured into ice-water, stirred until solid formed. The solid was filtered to give 2,5-dimethyl-7-(2,4,6-trimethylphenyl)-5H-pyrrolo[3,2-d]pyrimidin-4-ol. The alternative way for the work up is via extraction technique. The ice water was extracted with chloroform. The organic layer was dried and concentrated to dryness to give the desired product.

<sup>1</sup>H NMR δ (CDCl<sub>3</sub>) 6.92(s,2H), 6.88(s,1H), 4.14(s,3H), 2.43(s,3H), 2.29(s,3H), 2.09(s,6H).

The following compounds can be prepared in a similar manner:

- 2,5-Dimethyl-7-(2,6-dimethyl-4-chloro-phenyl)-5H-pyrrolo[3,2-d]pyrimidin-4-ol;
- 2,5-Dimethyl-7-(2,6-dimethyl-4-bromo-phenyl)-5H-pyrrolo[3,2-d]pyrimidin-4-ol;
- 2-Methyl-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo[3,2-d]pyrimidin-4-ol;
- 2-Methyl-7-(2,6-dimethyl-4-chloro-phenyl)-5H-pyrrolo[3,2-d]pyrimidin-4-ol;
- 2-Methyl-7-(2,6-dimethyl-4-bromo-phenyl)-5H-pyrrolo[3,2-d]pyrimidin-4-ol; and
- 2-Methyl-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo[3,2-d]pyrimidin-4-ol;

### PREPARATION H

#### 4-Chloro-2,5-dimethyl-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo[3,2-d]pyrimidine

A mixture of 2,5-dimethyl-7-(2,4,6-trimethylphenyl)-5H-pyrrolo[3,2-d]pyrimidin-4-ol (1 mmol) in POCl<sub>3</sub> (1.3 ml) was heated at reflux until all the starting material were

-49-

consumed (about 1-3 hours). The mixture was concentrated to dryness. The residue was poured into ice-water and extracted with ethyl acetate. The organic layer was dried and concentrated to dryness to give the title compound.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.18(s,1H), 6.95(s,2H), 4.16(s,3H), 2.69(s,3H), 2.31(s,3H),  
5 2.05(s,6H) ppm.

#### PREPARATION I

##### 7-Amino-1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]pyridine-6-carboxylic acid methyl ester

A mixture of 3-amino-1-methyl-4-(2,4,6-trimethylphenyl)-1H-pyrrole-2-carbonitrile  
10 (2 mmol), methyl acetoacetate (4 mmol) and SnCl<sub>4</sub> (4 mmol) in 1, 2-dichloroethane was heated at reflux for about six hours until all the starting material were consumed. The mixture was quenched with acetone, basified with saturated NaHCO<sub>3</sub>, then filtered through Celite®. The filtrate was concentrated to dryness. The residue was quenched  
15 with water and extracted with chloroform. The chloroform layer was washed with brine, dried and concentrated to dryness to give 7-amino-1,5-dimethyl 3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]pyridine-6-carboxylic acid methyl ester. The product may be purified by trituration.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 6.91(s,2H), 6.85(s,1H), 6.28(brs,2H), 4.10(s,3H), 3.89(s,3H),  
2.62(s,3H), 2.28(s,3H), 2.09(s,6H) ppm.

20

#### PREPARATION J

##### 7-Bromo-1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-1H-pyrrolo[3,2-b]pyridine-6-carboxylic acid methyl ester


A mixture 7-amino-1, 5-dimethyl-3-(2, 4, 6-trimethylphenyl)-1H-pyrrolo[3, 2-  
b]pyridine-6-carboxylic acid methyl ester (1 mmol), n-butylnitrite (BuONO) (1.5 mmol)  
25 and CuBr<sub>2</sub> in acetonitrile was heated at 60-70°C until all starting material was consumed. The mixture was quenched with water and extracted with ethyl acetate. The organic layer was dried and concentrated to dryness to give the title compound.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.09(s,1H), 6.94(s,2H), 4.16(s,3H), 3.98(s,3H), 2.52(s,3H),  
2.30(s,3H), 2.04(s,6H) ppm. 2,7-dibromo-1,5-dimethyl-3-(2,4,6-trimethyl-phenyl)  
30 -1H-pyrrolo[3,2-b]pyridine-6-carboxylic acid methyl ester was also produced as a minor component. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.00(s,2H), 4.45(s,3H), 4.03(s,3H), 2.51(s,3H), 2.34(s,3H), 2.12(s,6H) ppm.

-50-

CLAIMS

1. A compound of the formula



10 or a pharmaceutically acceptable salt thereof, wherein the dashed lines represent optional double bonds;

A is nitrogen or CR<sup>7</sup>;

B is -NR<sup>1</sup>R<sup>2</sup>, -CR<sup>1</sup>R<sup>2</sup>R<sup>10</sup>-C(=CR<sup>2</sup>R<sup>11</sup>)R<sup>1</sup>, -NHCR<sup>1</sup>R<sup>2</sup>R<sup>10</sup>, -OCR<sup>1</sup>R<sup>2</sup>R<sup>10</sup>, -SCR<sup>1</sup>R<sup>2</sup>R<sup>10</sup>, -CR<sup>2</sup>R<sup>10</sup>NHR<sup>1</sup>, -CR<sup>2</sup>R<sup>10</sup>OR<sup>1</sup>, -CR<sup>2</sup>R<sup>10</sup>SR<sup>1</sup> or -COR<sup>2</sup>;

15 J and K are each independently nitrogen or carbon and both J and K are not nitrogens ;

D and E are each selected, independently, from nitrogen, CR<sup>4</sup>, C=O, C=S, sulfur, oxygen, CR<sup>4</sup>R<sup>6</sup> and NR<sup>8</sup> ;

G is nitrogen or carbon;

20 the ring containing D, E, G, K, and J in formula I may be a saturated or unsaturated 5-membered ring and may optionally contain one or two double bonds and may optionally contain from one to three heteroatoms in the ring and may optionally have one or two C=O or C=S groups;

25 R<sup>1</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one or two substituents independently selected from hydroxy, fluoro, chloro, bromo, iodo, -O-(C<sub>1</sub>-C<sub>4</sub> alkyl), CF<sub>3</sub>, -C(=O)O-(C<sub>1</sub>-C<sub>4</sub> alkyl), -OC(=O)(C<sub>1</sub>-C<sub>4</sub> alkyl), -OC(=O)N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -NHCO(C<sub>1</sub>-C<sub>4</sub> alkyl), -COOH, -COO(C<sub>1</sub>-C<sub>4</sub> alkyl), -CONH(C<sub>1</sub>-C<sub>4</sub> alkyl), -CON(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -S(C<sub>1</sub>-C<sub>4</sub> alkyl), -CN, -NO<sub>2</sub>, -SO(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>NH(C<sub>1</sub>-C<sub>4</sub> alkyl) and -SO<sub>2</sub>N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), wherein each of the C<sub>1</sub>-C<sub>4</sub> alkyl groups in the foregoing R<sup>1</sup> groups may optionally contain one or two double or triple bonds;

30 R<sup>2</sup> is C<sub>1</sub>-C<sub>12</sub> alkyl which may optionally contain from one to three double or triple bonds, aryl or (C<sub>1</sub>-C<sub>4</sub> alkylene)aryl, wherein said aryl and the aryl moiety of said (C<sub>1</sub>-C<sub>4</sub>

alkylene)aryl is selected from phenyl, naphthyl, thienyl, benzothienyl, pyridyl, quinolyl, pyrazinyl, pyrimidinyl, imidazolyl, furanyl, benzofuranyl, benzothiazolyl, isothiazolyl, pyrazolyl, pyrrolyl, indolyl, pyrrolopyridyl, oxazolyl and benzoxazolyl; C<sub>3</sub>-C<sub>8</sub> cycloalkyl or (C<sub>1</sub>-C<sub>6</sub> alkylene)(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), wherein one or two of the carbon atoms of said  
 5 cycloalkyl and the 5 to 8 membered cycloalkyl moieties of said (C<sub>1</sub>-C<sub>6</sub> alkylene)(C<sub>3</sub>-C<sub>8</sub> cycloalkyl) may optionally and independently be replaced by an oxygen or sulfur atom or by NZ<sup>2</sup> wherein Z<sup>2</sup> is selected from hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, benzyl and C<sub>1</sub>-C<sub>4</sub> alkanoyl, and wherein each of the foregoing R<sup>2</sup> groups may optionally be substituted with from one to three substituents independently selected from chloro, fluoro, hydroxy and C<sub>1</sub>-C<sub>4</sub>  
 10 alkyl, or with one substituent selected from bromo, iodo, C<sub>1</sub>-C<sub>6</sub> alkoxy, -OC(=O)(C<sub>1</sub>-C<sub>6</sub> alkyl), -OC(=O)N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -S(C<sub>1</sub>-C<sub>6</sub> alkyl), amino, -NH(C<sub>1</sub>-C<sub>2</sub> alkyl), -N(C<sub>1</sub>-C<sub>2</sub> alkyl)(C<sub>1</sub>-C<sub>4</sub> alkyl), -N(C<sub>1</sub>-C<sub>4</sub> alkyl)-CO-(C<sub>1</sub>-C<sub>4</sub> alkyl), -NHCO(C<sub>1</sub>-C<sub>4</sub> alkyl), -COOH, -COO(C<sub>1</sub>-C<sub>4</sub> alkyl), -CONH(C<sub>1</sub>-C<sub>4</sub> alkyl), -CON(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -SH, -CN, -NO<sub>2</sub>, -SO(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>NH(C<sub>1</sub>-C<sub>4</sub> alkyl) and -SO<sub>2</sub>N(C<sub>1</sub>-C<sub>4</sub>  
 15 alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl);

-NR<sup>1</sup>R<sup>2</sup> or CR<sup>1</sup>R<sup>2</sup>R<sup>10</sup> may form a saturated 3 to 8 membered carbocyclic ring which may optionally contain from one to three double bonds and wherein one or two of the ring carbon atoms of such 5 to 8 membered rings may optionally and independently be replaced by an oxygen or sulfur atom or by NZ<sup>3</sup> wherein Z<sup>3</sup> is  
 20 hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, benzyl or C<sub>1</sub>-C<sub>4</sub> alkanoyl;

R<sup>3</sup> is hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, -O(C<sub>1</sub>-C<sub>4</sub> alkyl), chloro, fluoro, bromo, iodo, (C<sub>1</sub>-C<sub>2</sub> alkylene)-O-(C<sub>1</sub>-C<sub>2</sub> alkyl), (C<sub>1</sub>-C<sub>2</sub> alkylene)-OH, or -S(C<sub>1</sub>-C<sub>4</sub> alkyl);

each R<sup>4</sup> is, independently, hydrogen, (C<sub>1</sub>-C<sub>6</sub> alkyl), fluoro, chloro, bromo, iodo, hydroxy, cyano, amino, (C<sub>1</sub>-C<sub>2</sub> alkylene)-OH, CF<sub>3</sub>, CH<sub>2</sub>SCH<sub>3</sub>, nitro, -O(C<sub>1</sub>-C<sub>4</sub> alkyl),  
 25 -N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -S(C<sub>1</sub>-C<sub>4</sub> alkyl), -CO(C<sub>1</sub>-C<sub>4</sub> alkyl), -C(=O)H or -C(=O)O(C<sub>1</sub>-C<sub>4</sub>alkyl);

R<sup>6</sup> is hydrogen, methyl or ethyl;

R<sup>8</sup> is hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl;

R<sup>5</sup> is phenyl, pyridyl, pyrazinyl, pyrimidyl, pyridazinyl and wherein each of the  
 30 foregoing R<sup>5</sup> groups is substituted with from one to four substituents R<sup>13</sup> wherein one to three of said substituents may be selected, independently, from fluoro, chloro, C<sub>1</sub>-C<sub>6</sub> alkyl and -O(C<sub>1</sub>-C<sub>6</sub> alkyl) and one of said substituents may be selected from bromo, iodo, formyl, OH, (C<sub>1</sub>-C<sub>4</sub> alkylene)-OH, (C<sub>1</sub>-C<sub>4</sub>alkylene)-O-(C<sub>1</sub>-C<sub>2</sub> alkyl), -CN, -CF<sub>3</sub>, -NO<sub>2</sub>,

-NH<sub>2</sub>, -NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -N(C<sub>1</sub>-C<sub>2</sub> alkyl)(C<sub>1</sub>-C<sub>6</sub> alkyl), -OCO(C<sub>1</sub>-C<sub>4</sub> alkyl), (C<sub>1</sub>-C<sub>4</sub> alkylene)-O-(C<sub>1</sub>-C<sub>4</sub> alkyl), -S(C<sub>1</sub>-C<sub>6</sub> alkyl), (C<sub>1</sub>-C<sub>4</sub> alkylene)-S-(C<sub>1</sub>-C<sub>4</sub> alkyl), -C(=O)O(C<sub>1</sub>-C<sub>4</sub> alkyl), -C(=O)(C<sub>1</sub>-C<sub>4</sub> alkyl), -COOH, -SO<sub>2</sub>NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>N(C<sub>1</sub>-C<sub>2</sub> alkyl)(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>NH<sub>2</sub>, -NHSO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl), -S(C<sub>1</sub>-C<sub>6</sub> alkyl) and -SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl), and wherein  
 5 each of the C<sub>1</sub>-C<sub>4</sub> alkyl and C<sub>1</sub>-C<sub>6</sub> alkyl moieties in the foregoing R<sup>5</sup> groups may optionally have one or two double bonds;

R<sup>7</sup> is hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, halo (e.g., chloro, fluoro, iodo or bromo), hydroxy, -O(C<sub>1</sub>-C<sub>4</sub> alkyl), -C(=O)(C<sub>1</sub>-C<sub>4</sub> alkyl), -C(=O)O(C<sub>1</sub>-C<sub>4</sub> alkyl), -OCF<sub>3</sub>, -CF<sub>3</sub>, -CH<sub>2</sub>OH or -CH<sub>2</sub>O(C<sub>1</sub>-C<sub>2</sub> alkyl);

10 R<sup>10</sup> is hydrogen, hydroxy, methoxy or fluoro;

R<sup>11</sup> is hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl; and

with the proviso that: a) when both J and K are carbons and D is CR<sup>4</sup> and E is nitrogen, then G can not be nitrogen; (b) when both J and K are carbons and D and G are nitrogens, then E can not be CR<sup>4</sup> or C=O or C=S; (c) when both J and K are  
 15 carbons and D and E are carbons, then G can not be nitrogen; (d) when G is carbon, it must be double banded to E; and (e) in the ring containing J, K, D, E and G, there can not be two double bonds adjacent to each other;

and the pharmaceutically acceptable salts of such compounds.

2. Compounds according to claim 1 wherein B is -NR<sup>1</sup>R<sup>2</sup>, -NHCHR<sup>1</sup>R<sup>2</sup>,  
 20 -OCHR<sup>1</sup>R<sup>2</sup> and R<sup>1</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, which may optionally be substituted with one fluoro, or C<sub>1</sub>-C<sub>4</sub> alkoxy group and which may optionally contain one double or triple bond; and R<sup>2</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl or (C<sub>1</sub>-C<sub>2</sub> alkyl)-CO-(C<sub>1</sub>-C<sub>2</sub> alkyl) which may optionally contain one double or triple bond.

3. Compounds according to claim 1, wherein B is -CHR<sup>1</sup>R<sup>2</sup>, -NR<sup>1</sup>R<sup>2</sup>,  
 25 -NHCHR<sup>1</sup>R<sup>2</sup>, -OCHR<sup>1</sup>R<sup>2</sup>, -SCHR<sup>1</sup>R<sup>2</sup>; and R<sup>1</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, which may optionally be substituted with one hydroxy, fluoro, CF<sub>3</sub>, cyclopropyl or C<sub>1</sub>-C<sub>4</sub> alkoxy group and which may optionally contain one double or triple bond; and R<sup>2</sup> is benzyl or C<sub>1</sub>-C<sub>6</sub> alkyl, which may optionally contain one double or triple bond, wherein said C<sub>1</sub>-C<sub>6</sub> alkyl and the phenyl moiety of said benzyl may optionally be substituted with one fluoro, hydroxy,  
 30 CF<sub>3</sub>, cyclopropyl, C<sub>1</sub>-C<sub>2</sub> alkyl, C<sub>1</sub>-C<sub>2</sub> alkoxy or chloro group.

4. Compounds according to claim 1 wherein R<sup>3</sup> is methyl.

5. Compounds according to claim 1 wherein R<sup>4</sup>, R<sup>6</sup>, R<sup>8</sup>, R<sup>9</sup>, and R<sup>12</sup> are hydrogen or methyl.

6. Compounds according to claim 1 wherein  $R^5$  is di- or tri-substituted phenyl in which the two or three substituents are independently selected from  $C_1-C_4$  alkyl,  $O-(C_1-C_4 \text{ alkyl})$ ,  $(C_1-C_4 \text{ alkylene})-O-(C_1-C_4 \text{ alkyl})$ ,  $CF_3$ ,  $OCF_3$ ,  $CHO$ ,  $(C_1-C_4 \text{ alkylene})-OH$ , cyano, chloro, fluoro, bromo and iodo, wherein each of the  
5 forgoing  $(C_1-C_4)$  alkyl groups may optionally contain one double or triple bond.

7. Compounds 1 wherein  $R^3$  is methyl, ethyl, chloro or methoxy; and each of  $R^4$ ,  $R^6$ ,  $R^8$ ,  $R^9$ , and  $R^{12}$  is, independently, hydrogen, methyl or ethyl.

8. Compounds wherein  $R^5$  is di- or tri-substituted pyridyl, or pyrimidyl in which the two or three substituents are independently selected from  $C_1-C_4$  alkyl,  
10  $O-(C_1-C_4 \text{ alkyl})$ ,  $(C_1-C_4 \text{ alkylene})-O-(C_1-C_4 \text{ alkyl})$ ,  $CF_3$ ,  $OCF_3$ ,  $CHO$ ,  $(C_1-C_4 \text{ alkylene})-OH$ , cyano, chloro, fluoro, bromo and iodo, wherein each of the forgoing  $(C_1-C_4)$  alkyl groups may optionally contain one double or triple bond.

9. Compounds according to claim 1 wherein A is N, CH or  $CCH_3$ .

10. Compounds according to claim 1 wherein A is CH, J and K are carbon  
15 and D, E, and G are nitrogen.

11. Compounds according to claim 1 wherein J and D are nitrogen, and K and G are carbon, and E is CH,  $CCH_3$  or  $CC_2H_5$ .

12. Compounds according to claim 1 wherein J and K are carbon, and  $D-EG$  is  $O-C(CH_3)=C$ ,  $O-CH=C$ ,  $S-C(CH_3)=C$ ,  $S-CH=C$ ,  $N(CH_3)-C(CH_3)=C$ ,  
20  $NHC(CH_3)=C$ ,  $NHC(CH_3CH_2)=C$ ,  $N(CH_3)-CH=C$ ,  $O-N=C$ ,  $S-N=C$ ,  $N(CH_3)-N=C$ ,  $O-CH_2N$  or  $S-CH_2N$ .

13. A compound according to claim 1 wherein B is  $-CHR^1R^2$ ,  $-NCHR^1R^2$  or  $-OCHR^1R^2$ , and the  $CHR^1R^2$  group of B is a cyclopentane ring, a tetrahydrofuran ring  
or a tetrahydrothienyl ring.

25 14. A compound according to claim 1 wherein the  $NR^1R^2$  group of B is a five membered saturated or unsaturated heterocyclic ring.

15. A compound according to claim 14 wherein the  $NR^1R^2$  is a pyrrole ring, a pyrrolidino ring, a thiazolidino ring or a morpholino ring.

16. A pharmaceutical composition for the treatment, prevention or inhibition  
30 of (a) a disorder the treatment of which can be effected or facilitated by antagonizing CRF, including but not limited to disorders induced or facilitated by CRF, or (b) a disorder selected from inflammatory disorders such as rheumatoid arthritis and osteoarthritis, pain, asthma, psoriasis and allergies; generalized anxiety disorder; panic;

-54-

phobias; obsessive-compulsive disorder; post-traumatic stress disorder; sleep disorders induced by stress; pain perception such as fibromyalgia; mood disorders such as depression, including major depression, single episode depression, recurrent, depression, child abuse induced depression, mood disorders associated with  
 5 premenstrual syndrome, and postpartum depression; dysthemia; bipolar disorders; cyclothymia; chronic fatigue syndrome; stress-induced headache; cancer; irritable bowel syndrome, Crohn's disease; spastic colon; post operative ileus; ulcer; diarrhea; stress-induced fever; human immunodeficiency virus (HIV) infections; neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease and Huntington's disease;  
 10 gastrointestinal diseases; eating disorders such as anorexia and bulimia nervosa; hemorrhagic stress; chemical dependencies and addictions (e.g., dependencies on alcohol, cocaine, heroin, benzodiazepines, or other drugs); drug and alcohol withdrawal symptoms; stress-induced psychotic episodes; euthyroid sick syndrome; syndrome of inappropriate antidiarrhetic hormone (ADH); obesity; infertility; head traumas; spinal  
 15 cord trauma; ischemic neuronal damage (e.g., cerebral ischemia such as cerebral hippocampal ischemia); excitotoxic neuronal damage; epilepsy; stroke; immune dysfunctions including stress induced immune dysfunctions (e.g., porcine stress syndrome, bovine shipping fever, equine paroxysmal fibrillation, and dysfunctions induced by confinement in chickens, sheering stress in sheep or human-animal  
 20 interaction related stress in dogs); muscular spasms; urinary incontinence; senile dementia of the Alzheimer's type; multiinfarct dementia; amyotrophic lateral sclerosis; hypertension; tachycardia; congestive heart failure; osteoporosis; premature birth; and hypoglycemia in a mammal, comprising an amount of a compound according to claim 1 that is effective in the treatment of such disorder, and a pharmaceutically acceptable  
 25 carrier.

17. A method for the treatment, prevention or inhibition of (a) a disorder the treatment of which can be effected or facilitated by antagonizing CRF, including but not limited to disorders induced or facilitated by CRF, or (b) a disorder selected from  
 30 inflammatory disorders such as rheumatoid arthritis and osteoarthritis, pain, asthma, psoriasis and allergies; generalized anxiety disorder; panic; phobias; obsessive-compulsive disorder; post-traumatic stress disorder; sleep disorders induced by stress; pain perception such as fibromyalgia; mood disorders such as depression, including major depression, single episode depression, recurrent premenstrual syndrome, and



-55-

postpartum depression; dysthemia; bipolar disorders; cyclothymia; chronic fatigue syndrome; stress-induced headache; cancer; irritable bowel syndrome, Crohn's disease; spastic colon; post operative ileus; ulcer; diarrhea; stress-induced fever; human immunodeficiency virus (HIV) infections; neurodegenerative diseases such as

5 Alzheimer's disease, Parkinson's disease and Huntington's disease; gastrointestinal diseases; eating disorders such as anorexia and bulimia nervosa; hemorrhagic stress; chemical dependencies and addictions (e.g., dependencies on alcohol, cocaine, heroin, benzodiazepines, or other drugs); drug and alcohol withdrawal symptoms; stress-induced psychotic episodes; euthyroid sick syndrome; syndrome of inappropriate

10 antidiarrhetic hormone (ADH); obesity; infertility; head traumas; spinal cord trauma; ischemic neuronal damage (e.g., cerebral ischemia such as cerebral hippocampal ischemia); excitotoxic neuronal damage; epilepsy; stroke; immune dysfunctions including stress induced immune dysfunctions (e.g., porcine stress syndrome, bovine shipping fever, equine paroxysmal fibrillation, and dysfunctions induced by confinement

15 in chickens, sheering stress in sheep or human-animal interaction related stress in dogs); muscular spasms; urinary incontinence; senile dementia of the Alzheimer's type; multiinfarct dementia; amyotrophic lateral sclerosis; hypertension; tachycardia; congestive heart failure; osteoporosis; premature birth; and hypoglycemia in a mammal, comprising administering to a subject in need of said treatment an amount of a

20 compound according to claim 1, that is effective in treating such disorder.

18. A method of treating or preventing a disorder or condition, the treatment or prevention of which can be effected or facilitated by inhibiting CRH binding protein in a mammal, comprising administering to said mammal a CRH binding protein inhibiting amount of a compound according to claim 1.

25 19. A pharmaceutical composition for treating or preventing a disorder or condition, the treatment or prevention of which can be effected or facilitated by inhibiting CRH binding protein in a mammal, comprising a CRH binding protein inhibiting amount of a compound according to claim 1 and a pharmaceutically acceptable carrier.

30 20. A compound according to claim 11 or 12 wherein A is N or CH, R<sup>3</sup> is methyl and each R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>9</sup> and R<sup>12</sup> is, independently, hydrogen or methyl.


21. A compound according to claim 20, wherein R<sup>5</sup> is di- or tri-substituted phenyl, wherein the two or three substituents are independently selected from C<sub>1</sub>-C<sub>4</sub>,

-56-

alkyl, O-(C<sub>1</sub>-C<sub>4</sub> alkyl), (C<sub>1</sub>-C<sub>4</sub> alkylene)-O-(C<sub>1</sub>-C<sub>4</sub>alkyl), CF<sub>3</sub>, OCF<sub>3</sub>, CHO, (C<sub>1</sub>-C<sub>4</sub>alkylene)-OH, cyano, chloro, fluoro, bromo and iodo, wherein each of the foregoing (C<sub>1</sub>-C<sub>4</sub>) alkyl groups may optionally contain one double or triple bond.

22. A compound of the formula

5




10

or

15

20



25 wherein R<sup>3'</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl, R<sup>7'</sup> is hydrogen, methyl, chloro, bromo, -COOH or -COO(C<sub>1</sub>-C<sub>4</sub> alkyl), T is chloro, bromo, iodo or triflate, R<sup>8</sup> is hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl and R<sup>4</sup> is hydrogen, (C<sub>1</sub>-C<sub>8</sub> alkyl), fluoro, chloro, bromo, iodo, hydroxy, cyano, amino, (C<sub>1</sub>-C<sub>2</sub> alkylene)-OH, CF<sub>3</sub>, CH<sub>2</sub>SCH<sub>3</sub>, nitro, -O(C<sub>1</sub>-C<sub>4</sub> alkyl), -N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -S(C<sub>1</sub>-C<sub>4</sub> alkyl), -CO(C<sub>1</sub>-C<sub>4</sub> alkyl), -C(=O)H or -C(=O)O(C<sub>1</sub>-C<sub>4</sub>alkyl);

30

23. A compound according to claim 1 wherein said compound is:

7-(1-ethyl-propoxy)-5-methyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidine;

[2,5-Dimethyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-7-yl]-(1-ethyl-

propyl)-amine;

-57-

(1-Ethyl-propyl)-[5-methyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-7-yl]-amine;

7-(1-Ethyl-propoxy)-2,5-dimethyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidine;

5 [2,5-Dimethyl-3-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-7-yl]-ethyl-propyl-amine;

[6-Bromo-5-bromomethyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-7-yl]-(1-ethyl-propyl)-amine;

10 (1-Ethyl-propyl)-[5-methyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-7-yl]-amine;

[6-Bromo-5-methyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-b]pyridin-7-yl]-(1-ethyl-propyl)-methyl-amine;

7-(1-Ethyl-propoxy)-5-methyl-3-(2,4,6-trimethyl-phenyl)-3H-[1,2,3]triazolo[4,5-b]pyridine;

15 4-(1-Ethyl-propoxy)-2,5-dimethyl-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo[3,2-d]pyrimidine;

(±)-2,5-Dimethyl-4-(tetrahydro-furan-3-yloxy)-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo-[3,2-d]pyrimidine;

20 2,5-Dimethyl-4-(S)-(tetrahydro-furan-3-yloxy)-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo-[3,2-d]pyrimidine;

2,5-Dimethyl-4-(1-propyl-butoxy)-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo[3,2-d]pyrimidine; or

4-sec-Butylsulfanyl-2,5-dimethyl-7-(2,4,6-trimethyl-phenyl)-5H-pyrrolo[3,2-d]pyrimidine;

25 or a pharmaceutically acceptable salt of such compound.

# INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/IB 97/00922

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D487/04 C07D471/04 A61K31/505 A61K31/435  
 //(C07D487/04,239:00,231:00),(C07D471/04,249:00,221:00),  
 (C07D487/04,239:00,209:00)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95 33750 A (PFIZER) 14 December 1995 see claims 1,13	1,16
P,X	WO 96 35689 A (NEUROGEN) 14 November 1996 see page 3, line 18 - line 25; claim 1; examples IE,II,ID	1,16,22

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

27 October 1997

Date of mailing of the international search report

04.11.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Alfaro Faus, I

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB 97/00922

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claim(s) 17 and 18  
is(are) directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/IB 97/00922

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9533750 A	14-12-95	AU 2453095 A	04-01-96
		BR 9502708 A	30-04-96
		EP 0764166 A	26-03-97
		FI 964894 A	05-12-96
		HU 75774 A	28-05-97
		NO 965237 A	06-02-97
-----			
WO 9635689 A	14-11-96	US 5644057 A	01-07-97
		AU 5679096 A	29-11-96
		CA 2194756 A	14-11-96
		EP 0770080 A	02-05-97
-----			